


QUESTION 4

$$(a) \sum_{k=2}^5 (-1)^k k = (-1)^2 \times 2 + (-1)^3 \times 3 + (-1)^4 \times 4 + (-1)^5 \times 5 = -2.$$



$$(i) \quad x^2 = 4y \\ y = \frac{x^2}{4} \\ \frac{dy}{dx} = \frac{x}{2}.$$

$$\text{When } x = 2p, \frac{dy}{dx} = \frac{2p}{2} = p.$$

$$\text{Equation of tangent AP is} \\ y - y_1 = m(x - x_1) \\ y - p^2 = p(x - 2p) \\ y = px - p^2$$

$$(ii) \text{ Equation of normal BP is} \\ y - p^2 = -\frac{1}{p}(x - 2p).$$

B lies on BP at $x = 0$.

$$\text{When } x = 0, \quad y = p^2 - \frac{1}{p}(-2p) \\ = p^2 + 2.$$

$$\therefore B \text{ is } (0, p^2 + 2).$$

$$(iii) \text{ Substitute } y = 0 \text{ in } \textcircled{1}: 0 = px - p^2 \\ x = p.$$

$$\therefore A \text{ is } (p, 0).$$

If $C(x, y)$ is the midpoint of $A(p, 0)$ and

$$B(0, p^2 + 2), \quad x = \frac{p+0}{2} \text{ and } y = \frac{0+(p^2+2)}{2}.$$

$$x = \frac{p}{2} \quad \text{---}\textcircled{2}$$

$$y = \frac{p^2+2}{2} \quad \text{---}\textcircled{3}$$

$$\text{From } \textcircled{2}, \quad p = 2x.$$

$$\text{Substitute in } \textcircled{3}: y = \frac{4x^2+2}{2} = 2x^2+1.$$

$$\text{But } p > 0, \quad \therefore x > 0.$$

$$\therefore \text{ Cartesian equation of locus of } C \\ \text{ is } y = 2x^2 + 1, \quad x > 0.$$

$$(c) (i) \int_1^2 \frac{dx}{x} = [\ln x]_1^2 \\ = \ln 2 - \ln 1 \\ = \ln 2.$$

$$(ii) \int_1^2 \frac{dx}{x} \div \frac{b-a}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right], \\ \text{where } f(x) = \frac{1}{x}, \quad a = 1, \quad b = 2. \\ = \frac{2-1}{6} \left[\frac{1}{1} + 4 \times \frac{1}{1.5} + \frac{1}{2} \right] \\ = \frac{25}{36} (= 0.694).$$

$$(iii) \quad \ln 2 \div \frac{25}{36} \\ 2 \div e^{\frac{25}{36}} \\ 2^{\frac{36}{25}} \div e \quad (\text{raising both sides to power } \frac{36}{25}) \\ \therefore e \div 2.7132 \dots \\ = 2.713 \text{ (3 dec. places).}$$

QUESTION 5

$$(a) \text{ Prove } (n+1)(n+2) \dots (2n-1)2n \\ = 2^n [1 \times 3 \times \dots \times (2n-1)]$$

$$\text{If } n = 1, \quad \text{LHS} = 1 + 1 = 2$$

$$\text{RHS} = 2^1 \times 1 = 2.$$

\therefore The statement is true for $n = 1$.

Assume statement is true for $n = k$, that is, assume $(k+1)(k+2) \dots (2k-1)2k$

$$= 2^k [1 \times 3 \times \dots \times (2k-1)]. \quad \text{---}\textcircled{1}$$

Hence prove statement is true for $n = k+1$, that is, prove

$$(k+2)(k+3) \dots (2k+1)(2k+2) \\ = 2^{k+1} [1 \times 3 \times \dots \times (2k+1)]. \quad \text{---}\textcircled{2}$$

Now LHS

$$= (k+2)(k+3) \dots (2k+1)(2k+2) \\ = \frac{(k+1)(k+2)(k+3) \dots (2k-1)2k(2k+1)(2k+2)}{k+1} \\ = \frac{2k}{k+1} [1 \times 3 \times \dots \times (2k-1)](2k+1)(2k+2), \text{ from } \textcircled{1} \\ = \frac{2k}{k+1} [1 \times 3 \times \dots \times (2k-1)](2k+1)2(k+1) \\ = 2^{k+1} [1 \times 3 \times \dots \times (2k-1)(2k+1)] \\ = \text{RHS.}$$

\therefore If the statement is true for $n = k$, it is also true for $n = k+1$. But it is true for $n = 1$.

\therefore It is true for $n = 1 + 1 = 2$ and so on, that is, it is true for all integers $n \geq 1$.

$$(b) f(x) = e^x - 1 - x$$

$$(i) \quad f'(x) = e^x - 1$$

$$= 0 \text{ only when } x = 0.$$

\therefore There is only one stationary point (at $x = 0$).

$$f''(x) = e^x > 0 \text{ for all } x.$$


\therefore The graph of $f(x)$ is concave up for all x .

Since $f(x)$ is continuous for all x (being made up of the sum and difference of continuous functions), the stationary point at $x = 0$ is both a local and absolute minimum.

$$(ii) \text{ When } x = 0, \quad f(x) = e^0 - 1 - 0 = 0.$$

\therefore The least value of $f(x) = 0$.

$\therefore f(x) \geq 0$ for all x .



N.B. The gradient of $y = e^x - 1$ at $x = 0$ is 1, so $y = x$ is a tangent at $(0, 0)$.

This is also implied by (ii).

$$(iv) \text{ Inverse relation of } y = e^x - 1 \text{ is } x = e^y - 1.$$

$$\text{That is, } e^y = x + 1$$

$$y = \log_e(x + 1)$$

$$\therefore g^{-1}(x) = \log_e(x + 1).$$

$$(v) \text{ Domain of } g^{-1}(x) \text{ is } x + 1 > 0, \\ \text{that is, } x > -1.$$

$$(vi) \quad g(x) = e^x - 1$$

$$g^{-1}(x) = \log_e(1+x).$$

The graphs of a pair of inverse functions are symmetrical about the line $y = x$.

The graph of $y = g(x)$ is above the graph of $y = x$ except at $x = 0$ where they coincide.

\therefore The graph of $y = g^{-1}(x)$ is below the graph of $y = x$ except at $x = 0$ where they coincide.

$$\therefore \log_e(1+x) \leq x \text{ for all } x > -1.$$

QUESTION 6

$$(a) x = \cos^2 3t, \quad t > 0.$$

(i) Substitute $x = \frac{3}{4}$ in $\textcircled{1}$:

$$\frac{3}{4} = \cos^2 3t$$

$$\cos 3t = \pm \frac{\sqrt{3}}{2}$$

$$3t = \frac{\pi}{6}, \dots$$

$$t = \frac{\pi}{18}, \dots$$

Particle is first at $x = \frac{3}{4}$ after $\frac{\pi}{18}$ seconds.

$$(ii) v = \frac{dx}{dt} = 2 \cos 3t \cdot -3 \sin 3t \\ = -3 \sin 6t.$$

$$\text{When } t = \frac{\pi}{18}, \quad v = -3 \sin \left(6 \times \frac{\pi}{18} \right)$$

$$= -3 \sin \frac{\pi}{3}$$

$$= -\frac{3\sqrt{3}}{2} < 0.$$

Since $v < 0$, the particle is travelling in the negative direction.

$$(iii) a = \frac{dv}{dt} = -3 \times 6 \cos 6t \\ = -18 \cos 6t.$$

$$\cos 6t = 2 \cos^2 3t - 1 \\ \text{(using } \cos 2x = 2 \cos^2 x - 1) \\ = 2x - 1, \text{ from } \textcircled{1}.$$

$$\therefore a = -18(2x - 1).$$

$$(iv) a = -18(2x - 1)$$

$$= -36 \left(x - \frac{1}{2} \right)$$

$$= -6^2 \left(x - \frac{1}{2} \right),$$


which is of the form $\ddot{x} = -n^2(x - b)$, indicating simple harmonic motion with centre of oscillation at $x = \frac{1}{2}$.

$$(v) \text{ Period} = \frac{2\pi}{n} \text{ seconds}$$

$$= \frac{2\pi}{6} \text{ seconds}$$

$$= \frac{\pi}{3} \text{ seconds.}$$

(b) (i)



$$OD = 10 \text{ cm (radius)}$$

$$\therefore FD = 8 \text{ cm (Pythagoras' theorem).}$$

- (ii) OBC is a sector of a circle, centre O , radius 10 cm.


$$l = r\theta$$

$$4 = 10 \times \angle BOC$$

$$\therefore \angle BOC = 0.4 \text{ radians}$$

$$\angle DFE = \angle BOC$$

$$\therefore \angle DFE = 0.4 \text{ radians.}$$




$$ED^2 = FD^2 + FE^2 - 2 \times FD \times FE \cos 0.4$$

(by cosine rule)

$$= 8^2 + 8^2 - 2 \times 8 \times 8 \cos 0.4$$

$$= 128(1 - \cos 0.4).$$

Alternatively:



$$\cos \angle EOD = \frac{OE^2 + OD^2 - ED^2}{2 \times OE \times OD}$$


$$= \frac{10^2 + 10^2 - 128(1 - \cos 0.4)}{2 \times 10 \times 10}$$

$$\angle DOE = 0.3192 \dots$$

$$\approx 0.319 \text{ radians (3 dec. places).}$$

QUESTION 7

- (a) Initial conditions for velocity:



When $t = 0$,

$$\dot{x} = 30 \cos(5^\circ), \quad \dot{y} = -30 \sin(5^\circ). \quad \text{--- ①}$$

- (i) $\ddot{x} = 0$

$$\therefore \dot{x} = C_1 \text{ (constant).}$$

$$\therefore \dot{x} = 30 \cos(5^\circ) \text{ from ①.} \quad \text{--- ②}$$

$$x = \int 30 \cos(5^\circ) dt$$

$$= 30t \cos(5^\circ) + C_2.$$

When $t = 0$, $x = 0$, $\therefore C_2 = 0$.

$$\therefore x = 30t \cos(5^\circ).$$

$$\ddot{y} = -10$$

$$\therefore \dot{y} = \int -10 dt$$

$$= -10t + D_1.$$

When $t = 0$, $\dot{y} = -30 \sin(5^\circ)$ from ①.

$$\therefore D_1 = -30 \sin(5^\circ)$$

$$\therefore \dot{y} = -10t - 30 \sin(5^\circ). \quad \text{--- ③}$$

$$y = \int -10t - 30 \sin(5^\circ) dt$$

$$= -5t^2 - 30t \sin(5^\circ) + D_2.$$

When $t = 0$, $y = 0$, $\therefore D_2 = 0$.

$$\therefore y = -30t \sin(5^\circ) - 5t^2. \quad \text{--- ④}$$

- (ii) Ball strikes the ground when $y = -2$.
Substitute $y = -2$ in ④:

$$-2 = -30t \sin 5^\circ - 5t^2$$

$$5t^2 + 30t \sin 5^\circ - 2 = 0$$

$$t = \frac{-30 \sin 5^\circ \pm \sqrt{(-30 \sin 5^\circ)^2 - 4 \times 5 \times (-2)}}{2 \times 5}$$

$$= \frac{-30 \sin 5^\circ \pm \sqrt{900 \sin^2 5^\circ + 40}}{10}$$

(other answer negative and therefore irrelevant)


$$= 0.4229 \dots$$

- \therefore The ball strikes the ground after 0.42 seconds (2 dec. places).

- (iii) When $t = 0.4229$,

$$\dot{x} = 30 \cos(5^\circ) \text{ from ②,}$$

$$\dot{y} = -4.229 - 30 \sin(5^\circ), \text{ from ③.}$$



$$\tan \theta = \frac{4.229 + 30 \sin 5^\circ}{30 \cos 5^\circ}$$

$$= 0.22899 \dots$$

$$\theta \approx 12.9^\circ.$$

Angle at which the ball strikes the ground is 13° (nearest degree).

(b) $(1-x)^n = \binom{n}{0} - \binom{n}{1}x + \binom{n}{2}x^2 - \dots + \binom{n}{n}(-1)^n x^n$

$$\left(1 + \frac{1}{x}\right)^n = \binom{n}{0}\left(\frac{1}{x}\right)^0 + \binom{n}{1}\left(\frac{1}{x}\right)^1 + \binom{n}{2}\left(\frac{1}{x}\right)^2 + \dots + \binom{n}{n}(-1)^n\left(\frac{1}{x}\right)^n.$$

The term in x^2 in $(1-x)^n\left(1 + \frac{1}{x}\right)^n$ is

$$\binom{n}{2}\binom{n}{0}x^2\left(\frac{1}{x}\right)^0 - \binom{n}{3}\binom{n}{1}x^3\left(\frac{1}{x}\right)^1 + \binom{n}{5}\binom{n}{3}x^5\left(\frac{1}{x}\right)^3 + \dots + (-1)^n\binom{n}{n}\binom{n}{n-2}x^n\left(\frac{1}{x}\right)^n.$$

\therefore The coefficient of x^2 in $(1-x)^n\left(1 + \frac{1}{x}\right)^n$ is

$$\binom{n}{2}\binom{n}{0} - \binom{n}{3}\binom{n}{1} + \dots + (-1)^n\binom{n}{n}\binom{n}{n-2},$$

and this is the expression given in the question.

$$\text{Now } (1-x)^n\left(1 + \frac{1}{x}\right)^n = \left[(1-x)\left(1 + \frac{1}{x}\right)\right]^n$$

$$= \left(\frac{1}{x} - x\right)^n.$$

The general term of $\left(\frac{1}{x} - x\right)^n$ is

$$\binom{n}{r}\left(\frac{1}{x}\right)^{n-r}(-x)^r = \binom{n}{r}(-1)^r x^{2r-n}.$$

The term in x^2 has $2r - n = 2$

$$r = \frac{n+2}{2}.$$

$$\therefore \text{The coefficient of } x^2 = \binom{n}{\frac{n+2}{2}}(-1)^{\frac{n+2}{2}},$$

and only exists if n is even,

$\left(\frac{n+2}{2}\right)$ must be an integer.

$$\therefore \binom{n}{2}\binom{n}{0} - \binom{n}{3}\binom{n}{1} + \dots + (-1)^n\binom{n}{n}\binom{n}{n-2}$$

$$= \begin{cases} \left(\frac{n+2}{2}\right)(-1)^{\frac{n+2}{2}} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

END OF 3 UNIT (ADDITIONAL) AND 3/4 UNIT (COMMON) MATHEMATICS SOLUTIONS