

Saint Ignatius' College Riverview

Mathematics Extension 2

Trial HSC Examination 2001

- (a) If $P = 2 - ti$ where t is real, find \overline{iP} .

(b) The complex number, u , is given by $u = \frac{\sqrt{3}-i}{1+i}$

Find (i) The modulus of u (ii) The exact value of $\arg u$ (iii) u^6 in the form $a + bi$

(c) On an Argand diagram shade the region satisfied by both of the conditions:
 $|z - 2| \geq 1$ and $-\frac{\pi}{6} \leq \arg z \leq \frac{\pi}{6}$.

(d) If the complex number $Q = \frac{z-1}{z-2i}$, is purely imaginary (where $z = x + iy$) determine the Cartesian equation for the locus of z and sketch this locus.
- (a) Find $\int \operatorname{cosec} \theta \, d\theta$ using $t = \tan \frac{\theta}{2}$

(b) Evaluate $\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \frac{dy}{1+\cos y}$

(c) Evaluate $\int_0^{\frac{\pi}{4}} e^x \sin^2 x \, dx$

(d) Find $\int \frac{t^2-t-21}{(t^2+4)(2t-1)} \, dt$
- (a) Consider the function $f(x) = 9 - x^2$. On three separate sets of axes, sketch the following, showing all important features.

(i) $y = f(x)$ (ii) $y = |f(x)|$ (iii) $|y| = f(x)$

(b) Consider the function $y = \sin(\cos^{-1} x)$

(i) Find the domain and range of the function.

(ii) Sketch this function showing the important features.

(c) Consider the function f and g defined by:
 $f(x) = \frac{x+1}{x-2}$ for $x \neq 2$ and $g(x) = [f(x)]^2$

(i) Sketch the hyperbola $y = f(x)$, clearly labelling the horizontal and vertical asymptotes and the points of intersection with the x and y axes.

(ii) Sketch the curve $y = g(x)$ on a separate diagram showing all important features including any turning points.

(iii) On a separate diagram sketch the curve given by $y = g(-x)$
- (a) (i) On the same number plane sketch the graphs of $y = |x| - 3$ and $y = 5 + 4x - x^2$

(ii) Hence, or otherwise, solve $\frac{|x|-3}{5+4x-x^2} > 0$

(b) Given the polynomial $Q(x)$, where $Q(x) = kx^{k+1} - (k+1)x^k + 1$ ($k \neq 0$) prove that $Q(x)$ is divisible by $(x-1)^2$.

(c) The equation $x^3 + 2x - 1 = 0$ has roots p, q and r . Find

(i) the value of $p^2 + q^2 + r^2$

(ii) the equation with roots $-p, -q$ and $-r$

(d) $P(x)$ is a polynomial with the following form:


$$P(x) = Kx^3 + Mx^2 + Lx + N \text{ where } K, M, L \text{ and } N \text{ are real.}$$

$P(x)$ has roots of 5 and i and when divided by $(x - 2)$ the remainder is 3. Find $P(x)$.

5. (a) (i) Show, using differentiation, that the equation of the normal to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ at the point $P(a \sec \alpha, b \tan \alpha)$ is $ax \tan \alpha + by \sec \alpha = (a^2 + b^2) \sec \alpha \tan \alpha$

(ii) The vertical line through P meets an asymptote of the above hyperbola at M . The normal at P meets the x axis at K . Show that KM is at right angles to the asymptote.

(b) The following diagram shows a circle with centre at the origin O . The point $E(0, a)$ is fixed where $a > 3$. Q lies on the circle such that the angle EQF is a right angle and $EQ = QF$.



(i) Copy the diagram.

(ii) Show by substitution that $Q(3 \cos \alpha, 3 \sin \alpha)$ satisfies $x^2 + y^2 = 9$

(iii) Prove, using congruent triangles or otherwise, that F has coordinates $(3 \cos \alpha + a - 3 \sin \alpha, 3 \cos \alpha + 3 \sin \alpha)$

(iv) Find the locus of F as Q moves on the circle.

(v) Prove that the locus of the circle is independent of the value of a .

6. (a) The base of a solid is the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ where $b > a$. Sections perpendicular to the y -axis are squares with one side in the base of the solid. Show that the volume of the solid is $\frac{16a^2b}{3}$ cubic units.

(b) The curve $y = \sin x$ is revolved about the straight line $y = 1$. Use a slicing technique to find the volume of the solid of revolution formed by the portion of the curve from $x = 0$ to $x = \frac{\pi}{2}$.

(c) The area enclosed by $y = (x-2)^2$ and the straight line $y = 4$ is rotated about the y -axis. Using the method of cylindrical shells, find the volume of the solid formed.

7. (a) A certain particle of unit mass moving through air experiences air resistance proportional to the square of its speed, v metres per second.

(i) Explain why the equations of motion with upwards taken as positive are:

$$\ddot{x} = -g - kv^2, \text{ when moving upwards and}$$

$$\ddot{x} = -g + kv^2, \text{ when moving downwards,}$$

where g is the acceleration due to gravity and k is a positive constant.

(ii) Suppose that the particle is fired vertically upwards from the ground with an initial speed of u metres per second.


(α) Show that the maximum height, H metres, reached by the particle is:

$$H = \frac{1}{2k} \ln\left(1 + \frac{ku^2}{g}\right)$$

(β) Show that the time, T seconds, taken to reach maximum height is:

$$T = \frac{1}{\sqrt{gk}} \tan^{-1}\left(\frac{u\sqrt{k}}{\sqrt{g}}\right)$$

(b) (i) A particle of mass m travels with constant velocity v in a horizontal circle of radius R , centre C , around a track banked at an angle α to the horizontal, as shown in the diagram.



Show that if there is no tendency for the particle to slip sideways then $v = \sqrt{Rg \tan \alpha}$.

(ii) A particle travels in a horizontal circle of radius 1 metre around the lower half of the track where the angle of banking is given by $\tan^{-1}(\frac{5}{18})$. Another particle travels in a horizontal circle of radius 1.2 metres around the upper half of the track where the angle of banking is given by $\tan^{-1}(\frac{16}{27})$. Each particle travels with constant velocity so that it has no tendency to slip sideways. The particles are initially observed to be alongside each other. Taking $g = 10$ metres per second squared, find the time that elapses before the particles are next observed to be alongside each other.

8. (a) Prove that:

(i) $p^2 + q^2 \geq 2pq$


(ii) $k^4 + l^4 + m^4 + n^4 \geq 4klmn$, if k, l, m and n are positive.

(b) Given $z = \cos \alpha + i \sin \alpha$, where $\sin \alpha \neq 0$:

(i) Prove that $\frac{1}{1-z \cos \alpha} = 1 + i \cot \alpha$.

(ii) Hence, by considering $\sum_{k=0}^{\infty} (z \cos \alpha)^k$, deduce the sum of the infinite series $\sin \alpha \cos \alpha + \sin 2\alpha \cos^2 \alpha + \cdots + \sin k\alpha \cos^k \alpha + \cdots$.

(c)



AB is a tangent to the circle at P . M and K move on PR and PQ respectively so that KM is parallel to AB . Prove that the point of intersection of the perpendicular bisectors of QK and RM moves on a straight line.