

ASCHAM SCHOOL
MATHEMATICS EXTENSION 2
TRIAL EXAMINATION

2003

Time : 3 hours + 5 minutes reading time

Instructions:

Attempt all questions

All questions are of equal value

All necessary working should be shown for every question.

Full marks may not be awarded for careless or badly arranged work

A Table of Standard Integrals is provided

Approved calculators may be used

Each question should be answered in a separate booklet

Question 1

- a) $(2-3i)(4+i) = p+iq$ where $p, q \in R$. Find p and q . [1]
- b) (i) Express $z = -\sqrt{3} + i$ in modulus-argument form. [2]
- (ii) Hence show that $z^7 + 64z = 0$ [2]
- c) Sketch the following subsets of the Argand diagram, showing important features and intercepts with the axes.
- (i) $\{z : 1 < |z| \leq 3 \text{ and } 0 < \arg z \leq \frac{\pi}{2}\}$ [2]
- (ii) $\{z : |z+1| + |z-1| = 3\}$ [3]
- (iii) $\{z : \arg(z-2) - \arg(z+2) = \frac{\pi}{3}\}$ [2]
- d) Find the Cartesian form of the equation of the locus of the point z if $\operatorname{Re}\left[\frac{z-4}{z}\right] = 0$ [3]


Question 2 *Please take a new booklet*

- a) Find $\int \frac{e^{2x}}{e^x + 1} dx$ [2]
- b) Evaluate $\int \tan^3 x dx$ [2]
- c) Evaluate $\int_0^{\pi} e^x \sin x dx$ [3]
- d) (i) Show that $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{x(\pi-2x)} = \frac{2}{\pi} \ln 2$ [3]
- (ii) Using the substitution $u = a+b-x$, show that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$ [2]
- (iii) Evaluate $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos^2 x}{x(\pi-2x)} dx$ [3]

Question 3 *Please take a new booklet*

- a) A chocolate has a circular base of radius 1 cm. If every section perpendicular to this base is an equilateral triangle, find the volume of chocolate needed to make a box of 40 such chocolates. [6]

b)



The arch $y = \sin x$, $0 \leq x \leq \pi$, is revolved around the line $y = c$ to generate the solid shown.

- (i) Show that the volume generated is given by $\pi(\pi c^2 - 4c + \frac{\pi}{2})$ [6]
- (ii) Find the value of c which minimises the volume. [3]

Question 4 *Please take a new booklet*

- a) A ball of mass m is thrown vertically upwards under gravity, the air resistance to the motion being $\frac{mgv^2}{a^2}$ where the speed is v , a is a constant and g is the acceleration due to gravity.

- (i) Show that during the upward motion of the ball

$$v \frac{dv}{dx} = \frac{-g}{a^2} (a^2 + v^2)$$

where x is the upward displacement. [2]

- (ii) Show that the greatest height reached is $\frac{a^2}{2g} \ln \left(1 + \frac{u^2}{a^2} \right)$ where u is the speed of projection. [5]

- b) A curve is defined by the parametric equations $x = \cos^3 \theta$, $y = \sin^3 \theta$ for $0 < \theta < \frac{\pi}{4}$.
- (i) Show that the equation to the normal to the curve at the point $P (\cos^3 \phi, \sin^3 \phi)$ is $x \cos \phi - y \sin \phi = \cos 2\phi$ [4]
- (ii) The normal at P cuts the x -axis at A and the y -axis at B . Show $AB = 2\cot 2\phi$ [4]

Question 5 *Please take a new booklet*

- a) If $ax^3 + bx^2 + d = 0$ has a double root, show that $27a^2d + 4b^3 = 0$ [3]
- b) (i) Prove that $P(x) = \frac{1}{4}x^4 - \frac{1}{3}x^3 - 2x^2 + 4x + c$ has no real zeros if $c > 9\frac{1}{3}$ [4]
- (ii) Explain why the largest zero of $P(x)$ is greater than 2 if $c = -2$. Find an approximation for the largest zero of $P(x)$ using one application of Newton's method. [3]
- c) (i) P is any point inside a circle center O . M is the midpoint of chords AB through P . Find the locus of M . Explain your answer. [3]
- (ii) Q is any point outside a circle center C . N is the midpoint of chords DE through Q . State the locus of N . [2]


Continued on next page

Question 6 *Please take a new booklet*

- a) (i) Find the five fifth roots of unity. [2]
- (ii) If $\omega = \text{cis } \frac{2\pi}{5}$, show that $1 + \omega + \omega^2 + \omega^3 + \omega^4 = 0$ [3]
- (iii) Show that $z_1 = \omega + \omega^4$ and $z_2 = \omega^2 + \omega^3$ are roots of the equation $z^2 + z - 1 = 0$ [3]
- b) (i) By using the expansions of $\cos(x - y)$ and $\cos(x + y)$ show that $\sin x \sin y = \frac{1}{2}(\cos P - \cos Q)$ where $P = (x - y)$ and $Q = (x + y)$ [3]
- (ii) Hence prove that $\sin x + \sin 3x + \sin 5x + \dots + \sin(2n - 1)x = \frac{\sin^2 nx}{\sin x}$ [4]

Question 7 *Please take a new booklet*

- a) The graph of $y = x^3 + 3x^2 - 4$ is sketched below



- (i) Sketch the curves $y = |x^3 + 3x^2 - 4|$ and $y = \ln|x^3 + 3x^2 - 4|$ on separate axes. [3]
- (ii) Hence or otherwise determine the value of m , where m is a constant, such that the equation $2 \ln|x + 2| + \ln|x - 1| = m$ [4]

- b) AB is a diameter of a circle whose centre is O and C is a point on the circumference such that $\angle AOC$ is acute. OC is produced to meet the tangent at A in D. Let $\angle CBD = \alpha$ and $\angle ABC = \beta$. Prove

(i) $\tan(\alpha + \beta) = \frac{1}{2} \tan 2\beta$ [3]

(ii) $\tan \alpha = \tan^3 \beta$ [3]

(iii) Calculate the value of α when $AD = AB$ [2]

Question 8 *Please take a new booklet*

- a) (i) Show that the condition for the line $y = mx + c$ to be tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $c^2 = a^2 m^2 + b^2$ [3]

- (iii) Hence or otherwise prove that the pair of tangents from the point (3, 4) to the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$ are at right angles to each other. [4]

- b) Let $I_{2n} = \int_1^1 (1-x^2)^n dx$ where $n \geq 0$

(i) Use the substitution $x = \sin \theta$ to show that $I_{2n} = \frac{2n}{2n+1} I_{2n-2}$ [3]

(ii) Show that $I_6 = \frac{32}{35}$ [2]

(iii) Deduce that $I_{2n} = \frac{2^{2n+1} (n!)^2}{(2n+1)!}$ [3]

End of Examination