



**BAULKHAM HILLS HIGH SCHOOL**

**TRIAL 2014  
YEAR 12 TASK 4**

# Mathematics Extension 1

## General Instructions

- Reading time – 5 minutes
- Working time – 120 minutes
- Write using black or blue pen
- Board-approved calculators may be used
- Show all necessary working in Questions 11-14
- Marks may be deducted for careless or badly arranged work

**Total marks – 70**

**Exam consists of 11 pages.**

This paper consists of TWO sections.

**Section 1 – Page 2-4 (10 marks)**  
**Questions 1-10**

- Attempt Question 1-10

**Section II – Pages 5-10 (60 marks)**

- Attempt questions 11-14

**Table of Standard Integrals is on page 11**


**Section I - 10 marks**

Use the multiple choice answer sheet for question 1-10

1. Given the equation  $A = 10e^{-kt}$ , what is the value of  $k$  given that  $A = 3.6$  and  $t = 5$ .

- (A)  $-0.717$
- (B)  $-0.204$
- (C)  $0.204$
- (D)  $0.717$

2.



In the diagram above,  $TA$  is a tangent and  $PQ$  is a chord produced to  $T$ . The value of  $x$  is

- (A) 12
- (B)  $2\sqrt{3}$
- (C)  $4\sqrt{2}$
- (D)  $4\sqrt{6}$

3. How many distinct permutations of the letter of the word "DIVIDE" are possible in a straight line when the word begins and ends with the letter  $D$

- (A) 12
- (B) 180
- (C) 360
- (D) 720


4. The coordinates of the point that divides the interval joining  $(-7,5)$  and  $(-1, -7)$  externally in the ratio 1:3 are
- (A)  $(-10,8)$
- (B)  $(-10,11)$
- (C)  $(2,8)$
- (D)  $(2,11)$

5. What is the domain and range of  $y = 2 \cos^{-1} \frac{3x}{2}$  ?
- (A)  $D = \left\{x: -\frac{2}{3} \leq x \leq \frac{2}{3}\right\}, R = \{y: 0 \leq y \leq 2\pi\}$
- (B)  $D = \left\{x: -\frac{3}{2} \leq x \leq \frac{3}{2}\right\}, R = \{y: 0 \leq y \leq 2\pi\}$
- (C)  $D = \left\{x: -\frac{2}{3} \leq x \leq \frac{2}{3}\right\}, R = \left\{y: 0 \leq y \leq \frac{\pi}{2}\right\}$
- (D)  $D = \left\{x: -\frac{3}{2} \leq x \leq \frac{3}{2}\right\}, R = \left\{y: 0 \leq y \leq \frac{\pi}{2}\right\}$

6. Which of the following is the general solution of  $3 \tan^2 x - 1 = 0$ , where  $n$  is an integer?
- (A)  $n\pi \pm \frac{\pi}{6}$
- (B)  $n\pi \pm \frac{\pi}{3}$
- (C)  $2n\pi \pm \frac{\pi}{6}$
- (D)  $2n\pi \pm \frac{\pi}{3}$

7. The displacement of a particle moving in simple harmonic motion is given by  $x = 3 \cos \pi t$  where  $t$  is the time in seconds. The period of oscillation is:
- (A)  $\pi$
- (B)  $\frac{2\pi}{3}$
- (C) 2
- (D) 3

8.  $AB$  and  $CD$  are parallel chords in a circle, which are 10cm apart.  $OR \perp AB$ ,  $AB = 14\text{cm}$  and  $CD = 12\text{cm}$ .



Find the diameter of the circle to 1 decimal place

- (A) 4.4cm  
(B) 8.2cm  
(C) 14.8cm  
(D) 16.5cm
9. The domain of  $f(x) = \log_e[(x - 4)(5 - x)]$  is
- (A)  $4 \leq x \leq 5$   
(B)  $x \leq 4, x \geq 5$   
(C)  $4 < x < 5$   
(D)  $x < 4, x > 5$
10. Which of the following represents the derivate of  $y = \sin^{-1}\left(\frac{1}{x}\right)$ ?

(A)  $\frac{1}{x\sqrt{x^2-1}}$

(B)  $\frac{1}{\sqrt{x^2-1}}$

(C)  $\frac{-1}{x\sqrt{x^2-1}}$

(D)  $\frac{-1}{\sqrt{x^2-1}}$

**End of Section 1**

**Section II – Extended Response****All necessary working should be shown in every question.**

<b>Question 11 (15 marks) - Start on the appropriate page in your answer booklet</b>		<b>Marks</b>
a)	Evaluate $\int_0^{\frac{\pi}{4}} \cos^2 4x \, dx$	<b>3</b>
b)	Find $\int \frac{dx}{x(\log_e x)^{11}}$ , using the substitution $u = \log_e x$	<b>2</b>
c)	Prove the identity $\frac{1 + \sin 2x + \cos 2x}{\cos x + \sin x} = 2 \cos x$	<b>2</b>
d)	Solve for $x$ $\frac{4}{x-1} \leq 3$	<b>3</b>
e)	(i) Show that a root of the continuous function $f(x) = x^3 - \ln(x+1)$ lies between 0.8 and 0.9.	<b>1</b>
	(ii) Hence use the halving the interval method to find the value of the root correct to 1 decimal place.	<b>1</b>
f)	(i) Find $\frac{d}{dx} \left[ \tan^{-1} x + \tan^{-1} \left( \frac{1}{x} \right) \right]$	<b>2</b>
	(ii) Hence sketch $y = \tan^{-1} x + \tan^{-1} \left( \frac{1}{x} \right)$ for $-2 \leq x \leq 2$	<b>1</b>
<b>End of Question 11</b>		

**Question 12 (15 marks)** - Start on the appropriate page in your answer booklet


**Marks**

- a) When a polynomial  $P(x)$  is divided by  $x^2 - 4$  the remainder is  $2x + 3$ .  
What is the remainder when  $P(x)$  is divided by  $x - 2$

**2**

- b) In the given diagram,  $PQ$  and  $PR$  are tangents and  $Q, T, R$  are collinear.

**3**




Copy or trace the diagram in to your writing booklet.  
Prove that the points  $P, Q, S, R$  are concyclic.

Question 12 continues on the following page

Question 12 (continued)

c)



Not to Scale

Points  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  lies on the parabola  $x^2 = 4ay$ .  
The chord  $PQ$  subtends a right angle at the origin.

(i) Prove  $pq = -4$

2

(ii) Find the equation of the locus of  $M$ , the midpoint of  $PQ$ .

3

d) Find the coefficient of  $x^4$  in the expression of  $\left(x - \frac{2}{x}\right)^{12}$

2

e) Prove by mathematical induction

$$1 \times 2 + 2 \times 2^2 + 3 \times 2^3 + \dots + n \times 2^n = (n - 1)2^{n+1} + 2$$

for positive integers  $n \geq 1$

3


End of Question 12

<b>Question 13 (15 marks)</b> - Start on the appropriate page in your answer booklet		<b>Marks</b>
a)	(i) Express $\sqrt{3} \sin x - \cos x$ in the form $R \sin(x - \alpha)$ where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$ .	<b>2</b>
	(ii) Hence state the least value of $\sqrt{3} \sin x - \cos x$ and the smallest positive value of $x$ for this least value to occur.	<b>2</b>
b)	In the cubic equation $3x^3 - (2k - 4)x^2 + 5x + k^2 = 0$ the sum of the roots is equal to twice their product. Find the values of $k$ .	<b>3</b>
c)	Find the number of arrangements of the letters of the word <i>PENCILS</i> if there are 3 letters between <i>E</i> and <i>I</i> .	<b>2</b>
d)	Below is the graph of a function $y = f(x)$ <div style="text-align: center;"> </div> <p>Copy the diagram in your booklet, and on the same set of axes sketch a possible graph for <math>y = f'(x)</math>.</p>	<b>2</b>
e)	It is estimated that the rate of increase in the population of a particular species of bird is given by the equation $\frac{dP}{dt} = kP(L - P)$ where $k$ and $L$ are positive constants. <p>(i) Verify that for any positive constant <math>c</math>, the expression <math display="block">P = \frac{Lc}{c + e^{-kLt}}</math> satisfies the above differential equation.</p> <p>(ii) What can be deduced about <math>P</math> as <math>t</math> increases?</p>	<b>3</b> <b>1</b>
<b>End of Question 13</b>		



**Question 14 (15 marks)** - Start on the appropriate page in your answer booklet

a)



Water is poured into a conical vessel at a constant rate of  $24\text{cm}^3/\text{s}$ .  
The depth of water is  $h\text{cm}$  at any time  $t$  seconds.

- (i) Show that the volume of water is given by  $V = \frac{1}{3}\pi h^3$ . **1**
- (ii) Find the rate at which the depth of water is increasing when  $h = 16\text{cm}$ . **2**
- (iii) Hence find that rate of increase of the area of surface of the liquid when  $h = 16$ . **1**

b) The acceleration of a particle is given by the equation  $\frac{d^2x}{dt^2} = 8x(x^2 + 1)$ , where  $x$  is the displacement in centimetres from a fixed point  $O$ , after  $t$  seconds.  
Initially the particle is moving from  $O$  with speed  $2\text{cm/s}$  in a negative direction.

- (i) Prove the general result  $\ddot{x} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$ . **2**
- (ii) Hence show that the speed is given by  $2(x^2 + 1)\text{cm/s}$ . **2**
- (iii) Find an expression for  $x$  in terms of  $t$ . **2**

**Question 14 continues on the following page**

**Question 14 (continued)**

- c) A projectile is fired from the origin with velocity  $V$  with an angle of elevation  $\theta$ , where  $\theta \neq \frac{\pi}{2}$ .

YOU MAY ASSUME

$$x = Vt \cos \theta, \quad y = -\frac{1}{2}gt^2 + Vt \sin \theta$$

Where  $x$  and  $y$  are the horizontal and vertical displacements from  $O$ ,  $t$  seconds after firing

- (i) Show the equation of flight can be expressed as

$$y = x \tan \theta - \frac{1}{4h}x^2(1 + \tan^2 \theta) \quad \text{where } h = \frac{V^2}{2g} \quad \mathbf{2}$$

- (ii) Show that a point  $(X, Y)$  can be hit by firing at 2 different angles  $\theta_1$  and  $\theta_2$  provided  $X^2 < 4h(h - Y)$ .  $\mathbf{2}$

- (iv) Show that no point above the  $x$ -axis can be hit by firing at 2 different angles  $\theta_1$  and  $\theta_2$  satisfying both  $\theta_1 < \frac{\pi}{4}$  and  $\theta_2 < \frac{\pi}{4}$ .  $\mathbf{1}$

**End of Paper.**

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \quad \text{if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$


NOTE:  $\ln x = \log_e x$ ,  $x > 0$

# EXT 1 SOLUTIONS

1)  $3 \cdot 6 = 10 e^{-5k}$   
 $0.36 = e^{5k}$   
 $k = 0.204$  C

2)  $x^2 = 12 \times 8$   
 $= 96$   
 $x = \sqrt{96}$   
 $= 4\sqrt{6}$  D

3) D  $\frac{4!}{2!}$  D  
 $\frac{4 \times 3 \times 2}{2} = 12$   
A




$X = \frac{1x-1 + 3x-7}{-2}$   
 $= \frac{-1+21}{-2}$  B  
 $= -10$   
 $Y = \frac{1x-7 + 3x+5}{-2}$   
 $= \frac{-7+15}{-2} = 11$

5)  $y = 2 \cos^{-1} \frac{3x}{2}$   
 $\frac{y}{2} = \cos^{-1} \frac{3x}{2}$   
 $-1 \leq \frac{3x}{2} \leq 1$   
 $-\frac{2}{3} \leq x \leq \frac{2}{3}$   
 $0 \leq \frac{y}{2} \leq \pi$   
 $0 \leq y \leq 2\pi$  A

6)  $3 \tan^2 x - 1 = 0$   
 $\tan^2 x = \frac{1}{3}$   
 $\tan x = \pm \frac{1}{\sqrt{3}}$   
 $x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$   
 $2\pi - \frac{\pi}{6}, 2\pi + \frac{\pi}{6}$   
 $= n\pi \pm \frac{\pi}{6}$  A

7)  $x = 3 \cos \pi t$   
 Period =  $\frac{2\pi}{\pi}$   
 $= 2$   
C




$7^2 + x^2 = (10-x)^2 + 6^2$

$x = \frac{80}{27}$

$r = \sqrt{7^2 + \left(\frac{80}{27}\right)^2}$   
 $= 8.24 \dots$

$\therefore d = 16.5$  D

9)  $(x-4)(5-x) > 0$



$4 < x < 5$  C

10.  $y = \sin^{-1}\left(\frac{1}{x}\right)$   
 $\frac{dy}{dx} = \frac{1}{\sqrt{1-\left(\frac{1}{x}\right)^2}} \cdot \frac{-1}{x^2}$   
 $= \frac{-1}{x^2 \sqrt{1-\frac{1}{x^2}}}$   
 $= \frac{-1}{x \sqrt{x^2-1}}$  C

119) (i)  $f(x) = x^2 - \ln(x+1)$   
 $f(0.8) = -0.075$   
 $f(0.9) = 0.146$

$\therefore$  Root lies between 0.8 & 0.9.  $\therefore$  Next exists ①

Let 1st APPROX = 0.85.

$f(0.85) = -0.001$

$\therefore$  Root lies between 0.85 & 0.9 ①

$\therefore$  Root = 0.9 to 1 dec Place. ①

b)  $\int \frac{dx}{x(\log x)}$

$u = \log x$   
 $\frac{du}{dx} = \frac{1}{x} dx$  ①  
 $= \int \frac{du}{u^{11}}$   
 $= \int u^{-11}$   
 $= \frac{u^{-10}}{-10} + C$  ①  
 $= \frac{1}{-10} (\log x)^{-10} + C$

Q11

$\cos^2 2x = 2 \cos^2 x - 1$

$\cos^2 x = \frac{1}{2} [\cos 2x + 1]$

$\cos^2 4x = \frac{1}{2} [\cos 8x + 1]$

$\int_0^{\pi/4} \cos^2 4x dx = \frac{1}{2} \int_0^{\pi/4} \cos 8x + 1 dx$  ①

$= \frac{1}{2} \left[ \frac{1}{8} \sin 8x + x \right]_0^{\pi/4}$  ①

$= \frac{1}{2} \left[ \left(0 + \frac{\pi}{4}\right) - 0 \right]$

$= \frac{\pi}{8}$  ①

c) LHS =  $\frac{1 + 2 \sin x \cos x + 2 \cos^2 x - 1}{\cos x + \sin x}$  ①

$= \frac{2 \cos x + \sin x}{\cos x + \sin x}$  ①


$= \frac{2 \cos x (\sin x + \cos x)}{\cos x + \sin x}$

$= 2 \cos x$   
 $= RHS$

d)  $\frac{4}{x-1} \leq 3$

$4 \leq 3x-3$

$\frac{7}{3} \leq x$



$x < 1, x \geq 2\frac{1}{3}$

if  $x \leq 1, x \geq 2\frac{1}{3}$  2 marks

if  $1 < x \leq 2\frac{1}{3}$  2 marks


e)  $\frac{d}{dx} \left[ \tan^{-1} x + \tan^{-1} \left( \frac{1}{x} \right) \right]$

$= \frac{1}{1+x^2} + \frac{1}{1+(\frac{1}{x})^2} \cdot \left(-\frac{1}{x^2}\right)$

$= \frac{1}{1+x^2} - \frac{1}{x^2+1}$

$= 0$  ①

$\therefore y = \tan^{-1} x + \tan^{-1} \left( \frac{1}{x} \right)$  is horizontal



12a)  $P(x) = (x^2 - 4)(Q(x)) + 2x + 3$   
 $P(x) = 4x + 3$   
 $= 7$   
 $\therefore \text{Rem} = 7$

b)  $\angle \text{PRT} = \angle \text{ISR} = x^\circ$  (L between tang & chort)  
 $\text{①} = \text{L in Alt segment}$

SIMILARLY  
 $\angle \text{QAR} = \angle \text{OST} = y^\circ$

$\angle \text{QSR} = x^\circ + y^\circ$

$\angle \text{QPR} = 180 - x - y$  (sum of A)  
 $\text{①}$

$\therefore \angle \text{QSR} + \angle \text{QPR} = 180^\circ$   $\text{①}$

$\therefore P, Q, S, R$  are concyclic.

c) (i)  $m\angle \text{OP} = \frac{90^\circ - 0}{2 \times 90 - 0}$   
 $= \frac{90}{180} = \frac{1}{2}$   $\text{①}$

Similarly  $m\angle \text{OQ} = \frac{90}{180} = \frac{1}{2}$   
 $m\angle \text{OP} \times m\angle \text{OQ} = -1$  since they are  $\perp$ .

$\therefore \angle \text{P} \times \angle \text{Q} = -1$   $\text{①}$   
 $\therefore \angle \text{P} = -4$   $\text{①}$

(ii) Midpoint  $(\frac{2x+2y}{2}, \frac{y^2+xy}{2})$   
 $= (\frac{x+y}{2}, \frac{y^2+xy}{2})$   $\text{①}$   
 $x = a + (p+q)$   
 $\therefore p+q = \frac{x-a}{a}$   $y = a \frac{[(p+q)^2 - 2pq]}{2}$   $\text{①}$   
 $y = a \left[ \frac{x^2}{a^2} - 2(-1) \right]$   
 $y = \frac{x^2}{a} + 4a$   $\text{①}$  or equivalent for  $n=k$ .

12d)  $T_{k+1} = C_k^k x^k \left(\frac{-2}{x}\right)^{12-k}$   $\text{①}$   
 $= C_k^k x^k (-2)^{12-k} x^{k-12}$   
 $= C_k^k (-2)^{12-k} x^{2k-12}$   
 $\therefore 2k - 12 = 4$   
 $k = 8$   
 $\therefore \text{coeff} = C_8^8 (-2)^4 = 16$   $\text{①}$

e) Prove True  $N=1$  2 tickets - 1 mark  
 In  $1 \times 2 = (1-1)2^2 + 2$  3 tickets - 2 marks  
 $= 2$  4 tickets - 3 marks  
 Yes.  $\text{①}$

ASSUME TRUE  $N=k$ .

$1 \times 2 + 2 \times 2^2 + 3 \times 2^3 + \dots + k \times 2^k = (k-1)2^{k+1} + 2$

Prove True  $N=k+1$ .

$2 + \dots + k \times 2^k + \frac{(k+1) \times 2^{k+1}}{2} = k \cdot 2^{k+2} + 2$   $\text{①}$

LHS  $= (k-1)2^{k+1} + 2 + (k+1) \times 2^{k+1}$   
 $= 2^{k+1}(2k) + 2$

$= 2^{k+1} \times k + 2$

$= 2^{k+2} \times k + 2$   $\text{①}$

RHS. true for  $n=k+1$  if true for  $n=k$ . true for  $n=1, n=2 \rightarrow$  for all  $n \in \mathbb{N}$   $\text{①}$

13. a) (i)  $\sqrt{3} \sin x - \cos x = R \sin(x+d)$   
 $= R \sin x \cos d + R \cos x \sin d$

$\therefore R \cos d = \sqrt{3}$   
 $R \sin d = -1$

$\therefore \tan d = \frac{-1}{\sqrt{3}}$   
 $\therefore d = \frac{5\pi}{6}, \frac{11\pi}{6}$  (1)

but  $R > 0$ ,  $\cos d > 0$ ,  $\sin d < 0$   
 $\therefore d$  in Quad 4.  $\therefore d = \frac{5\pi}{6}$

$R^2 = (-1)^2 + (\sqrt{3})^2$   
 $= 4$   
 $R = 2$  (1)

$\therefore \sqrt{3} \sin x - \cos x = 2 \sin(x + \frac{5\pi}{6})$

(ii)  $\therefore$  Least value of  $\sqrt{3} \sin x - \cos x = -2$  (1)

$\sin(x + \frac{11\pi}{6}) = -1$

$x + \frac{11\pi}{6} = \frac{3\pi}{2}, \frac{7\pi}{2}$

but  $x > 0$   $x = \frac{2\pi}{6} - \frac{11\pi}{6} \Rightarrow x = \frac{5\pi}{3}$  (1)

13b)

$\alpha + \beta + \gamma = \frac{2K-4}{3}$

$\alpha \beta \gamma = -\frac{K^2}{3}$


$\therefore 2K-4 = -2K^2$  (1)

13c)

$E \dots I$   $2K^2 + 2K - 4 = 0$   
 $K^2 + K - 2 = 0$

(1) 3 letters between E & I in Brings 6  
 no. ways =  $6 \times 2 \times 2$  (4) correct  
 = 720 ways (1) paper  $\therefore K = -2, 1$  (1)

13d)



Correct  $0 \leq x \leq 2$  (1)

Correct the rest  
 $x < 0$   $x > 2$  (1)

e)  $\frac{dP}{dt} = kP(L-P)$

i)  $P = \frac{LC}{C + e^{-kLt}} = LC(C + e^{-kLt})^{-1}$

$\frac{dP}{dt} = -LC(C + e^{-kLt})^{-2} \cdot -kL e^{-kLt}$   
 $= \frac{kL^2 C e^{-kLt}}{(C + e^{-kLt})^2}$

CONSIDER:  $\frac{dP}{dt} = kP(L-P)$

$= k \cdot \frac{LC}{C + e^{-kLt}} \left( L - \frac{LC}{C + e^{-kLt}} \right)$   
 $= \frac{kLC}{(C + e^{-kLt})^2} \frac{LC + Le^{-kLt} - LC}{\dots}$

- (1) Differentiating
- (1) Sign. progress
- (1) fully correct.

$= \frac{kL^2 C e^{-kLt}}{(C + e^{-kLt})^2} = \frac{dP}{dt}$

ii) as  $t \rightarrow \infty$   $P \rightarrow L$  (1)

14a) by SIM Δ's

i)  $\frac{4r}{30} = \frac{h}{30}$  (matching sides in III Δ's)

$\therefore r = h$   
 $V = \frac{1}{3} \pi r^2 h$   
 $= \frac{1}{3} \pi h^2 \cdot h$   
 $= \frac{1}{3} \pi h^3$

ii)  $\frac{dV}{dt} = \frac{dh}{dt} \cdot \frac{dV}{dh}$   
 $24 = \frac{dh}{dt} \cdot \pi h^2$  ①  
 $h = 16$   
 $\frac{dh}{dt} = \frac{24 \cdot 3}{\pi \cdot 16^2}$   
 $= \frac{3}{32\pi} \text{ cm/s}$  ①

iii)  $S = \pi r^2$   
 $= \pi h^2$   
 $\frac{dS}{dt} = \frac{dS}{dh} \cdot \frac{dh}{dt}$   
 $= 2\pi h \cdot \frac{3}{32\pi}$  ①  
 $= 3 \text{ cm}^2/\text{s}$

b)  $\frac{d}{dx} \left( \frac{1}{2} v^2 \right) = \frac{d \left( \frac{1}{2} v^2 \right)}{dv} \cdot \frac{dv}{dx}$   
 $= v \frac{dv}{dx}$   
 $= \frac{dx}{dt} \cdot \frac{dv}{dx}$   
 $= \frac{dv}{dt}$   
 $= \dot{x}$

$\dot{x} = 8x(x^2+1)$

i)  $\frac{1}{2} v^2 = \int 8x^3 + 8x dx$

$\frac{1}{2} v^2 = \frac{8x^4}{4} + 4x^2 + C$  ①

$x=0, v=-2$   
 $-2 = C$

$\frac{1}{2} v^2 = 2x^4 + 4x^2 - 2$   
 $v^2 = 4x^4 + 8x^2 + 4$

$= 4(x^2 + 2x^2 + 1)$   
 $v^2 = 4(x^2 + 1)^2$

$v = \pm 2(x^2 + 1)$

speed =  $2(x^2 + 1) \text{ cm/s}$  ①

iii)  $v = \pm 2(x^2 + 1)$

but when  $x=0, v=-2$ .

$\therefore v = -2(x^2 + 1)$  ①

$\frac{dx}{dt} = -2(x^2 + 1)$

$\frac{dt}{dx} = \frac{1}{-2(x^2 + 1)}$

$t = -\frac{1}{2} \int \frac{1}{x^2 + 1} dx$

$t = -\frac{1}{2} \tan^{-1}(x) + C$

$t=0, x=0, \therefore C=0$

$t = -\frac{1}{2} \tan^{-1}(x)$

$-2t = \tan^{-1}(x) \implies x = \tan(-2t)$  ①  
 $= -\tan(2t)$

14c)  $x = vt \cos \theta$

$\therefore t = \frac{x}{v \cos \theta}$

$y = -\frac{1}{2} g \left( \frac{x}{v \cos \theta} \right)^2 + \frac{vx \sin \theta}{v \cos \theta}$  simplify to  
 $= -\frac{gx^2}{2v^2 \cos^2 \theta} + x \tan \theta$  ①

$= x \tan \theta - \frac{gx^2}{2v^2} \sec^2 \theta$

$= x \tan \theta - \frac{gx^2}{2v^2} (1 + \tan^2 \theta)$

but  $v^2 = 2gh$

$y = x \tan \theta - \frac{gx^2}{4gh} (1 + \tan^2 \theta)$  ①

$= x \tan \theta - \frac{x^2}{4h} (1 + \tan^2 \theta)$  ①

clearly show ① relationship

$\frac{gx^2}{4h}$



ii) If Point to Pass thru  $(X, Y)$

$$Y = X \tan \theta - \frac{X^2}{4h} (1 + \tan^2 \theta)$$

$$\therefore X^2 \tan^2 \theta - 4hX \tan \theta + (4hY + X^2) = 0 \quad \text{①}$$

FOR DIFFERENT ROOTS  $\Delta > 0$

$$\therefore 16h^2 X^2 - 4X^2(4hY + X^2) > 0 \quad \text{②}$$

$$4X^2(4h^2 - 4hY - X^2) > 0$$

$$\therefore \text{since } 4X^2 > 0, \quad 4h^2 - 4hY - X^2 > 0 \quad \text{③}$$

$$\therefore 4h(h - Y) > X^2 \quad \text{④}$$

Identifies tan  $\theta$  as the  
Variable of  $Y$  for

uses  $\Delta > 0$  to find  
2 solutions

ii) If  $\tan \theta_1, \tan \theta_2$  are roots of quadratic  
eqn

$$X^2 \tan^2 \theta - 4hX \tan \theta + (4hY + X^2) = 0$$

$$\therefore \tan \theta_1 \cdot \tan \theta_2 = \frac{4hY + X^2}{X^2}$$

$$= 1 + \frac{4hY}{X^2}$$

$$\therefore \tan \theta_1 \text{ or } \tan \theta_2 > 1$$

$$\therefore \theta_1 \text{ or } \theta_2 > \pi/4 \quad \text{①}$$