

**CRANBROOK**

**MATHEMATICS EXTENSION 2**

**2007**

**HIGHER SCHOOL CERTIFICATE**

**TRIAL EXAMINATION**

**General Instructions**

- Reading time – 5 minutes
- Writing time – 3 hours
- All eight questions should be attempted
- Total marks available - 120
- All questions are worth 15 marks
- An approved calculator may be used
- All relevant working should be shown for each question
- Answer each question in a separate 8 page booklet.

**Question 1** (15 marks)

Marked by SKB

**Marks**

(a) Find  $\int_1^2 3\sqrt{x-1} dx$  2

(b) Using the substitution  $u = e^x - 1$  or otherwise, 2

find  $\int \frac{2e^x}{e^{2x} - 2e^x + 1} dx$

(c) Use integration by parts to evaluate  $\int_0^{\frac{\pi}{4}} x \cos 4x dx$  3

(d) Use the substitution  $t = \tan \frac{\theta}{2}$  to find  $\int \frac{\sin \theta}{1 + \cos \theta} d\theta$  4

(e) (i) Find the real numbers  $a$ ,  $b$  and  $c$  such that 2

$$\frac{x^3 + 5x^2 + x + 2}{x^2(x^2 + 1)} \equiv \frac{x + a}{x^2} + \frac{bx + c}{x^2 + 1}$$

(ii) Find  $\int \frac{x^3 + 5x^2 + x + 2}{x^2(x^2 + 1)} dx$  2

**Question 2** (15 marks)

Marked by SKB

**Marks**

(a) Evaluate  $\int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\sin^3 x + \cos^3 x} dx$ . **2**

(b) Evaluate  $\int_0^1 \sqrt{1+x^2} dx$ . **4**

(c) Find  $\int \frac{3x-4}{\sqrt{4+5x-3x^2}} dx$  **4**

(d) **5**  
If  $I_n = \int_0^{\frac{\pi}{2}} \cos^n x \sin^2 x dx$  for  $n \geq 0$ , show that  
 $I_n = \frac{n-1}{n+2} I_{n-2}$  for  $n \geq 2$ . Hence or otherwise  
evaluate  $\int_0^{\frac{\pi}{2}} \cos^4 x \sin^2 x dx$ .

**Question 3** (15 marks)

Marked by CJL


**Marks**(a) Let  $z = 3 - 4i$  and  $\omega = 2 - i$ (i) Find  $\frac{1}{z}$  in the form  $x + yi$  **1**(ii) Show that  $\text{Im } z + \bar{\omega} + z\omega = -10i$  **2**(b) If  $ai$  is a solution to the equation **2**

$$z^2 + (1 - i)z + (2 - 2i) = 0$$

find the real value of  $a$ .(c) Let  $u = 1 - i$ (i) Find  $|u|$  and  $\arg u$  **2**(ii) Hence find  $u^{12}$ . Express your answer in the form  $x + yi$ . **2**(d) Sketch the region on an Argand diagram where the inequalities **3**

$$|z - 2 + i| \leq |z + 2 - i| \text{ and } \text{Im } z \geq 0 \text{ both hold.}$$

(e)



The point  $A$  on the Argand diagram above corresponds to the complex number  $z$ .

Triangle  $ABO$  is a right-angled triangle where  $OB = 2OA$ .

(i) Show that point  $B$  corresponds to the complex number  $2iz$ . **1**(ii) The point  $C$  corresponds to the complex number  $v$  and  $C$  is situated so that  $OACB$  is a rectangle.


Given that  $z = x + yi$ ,  $x, y \in \mathbb{R}$ , find  $\bar{v}$  in terms of  $x$  and  $y$ . **2**

**Question 4** (15 marks)      Marked by SKB

**Marks**

- (a) The base of a certain solid  $S$  lies on the ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$ . The cross-section of this solid by planes perpendicular to the  $x$ -axis are equilateral triangles. By including appropriate views of slices to this solid calculate its volume in exact form. 5


(b)



The shaded area shown in the diagram above is the area between the graph of  $y = 4x + 1$  and the graph of  $y = 2x^2 + 1$ . This shaded area is rotated about the  $y$  axis to form a solid.

Use the method of cylindrical shells to find the volume of the solid. 4

(c)



The shaded area is bounded by the lines  $x = -4, x = -2, y = 8$ , by the curve  $y = \frac{-8}{x+2}$  and by the  $x$ -axis.

The region is rotated about the line  $x = 3$  until it reaches its original position thus forming a solid. The horizontal line segment  $PQ$  forms an annulus as a result of this rotation.

- (i) Show that the area of this annulus at height  $y$  where  $y \geq 4$ , is equal to

$$16\pi \left( \frac{4}{y^2} + \frac{5}{y} \right) \quad 2$$

- (ii) Hence find the volume of the solid. 4

**Question 5** (15 marks)      Marked by CJL**Marks**


- (a) Let  $f(x) = \cos^{-1} x$  for  $-1 \leq x \leq 1$  and  $g(x) = \sin^{-1} x$  for  $-1 \leq x \leq 1$ .
- (i) Sketch  $f(x)$  and  $g(x)$  on the same set of axes. **1**
- (ii) By differentiating, evaluate  $f(x) + g(x)$  **1**
- (iii) Hence evaluate  $\int_{-1}^1 (f(x) + g(x)) dx$  **1**
- (b) The ellipse E has the equation  $x^2 + \frac{y^2}{4} = 1$ .
- (i) Find the eccentricity and the foci of E. **2**
- (ii) Find the length of the major and minor axes of E. **1**
- (iii) Write down the equations of the directrices of E. **1**
- (iv) Sketch E. **1**
- (c) (i) The polynomial equation  $p(x) = 0$  has a root  $\alpha$  of multiplicity 3. **2**  
Show that  $\alpha$  is a root of  $p'(x) = 0$  and is of multiplicity 2.
- (ii) The polynomial  $q(x) = x^6 + ax^5 + bx^4 - x^2 - 2x - 1$  has a quadratic factor of  $x^2 + 2x + 1$ . Find  $a$  and  $b$ . **2**
- (i) Consider the polynomial **3**  
$$r'(x) = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \text{ where } r(0) = 1.$$
 Show that  $r(x)$  has no double roots.

**Question 6** (15 marks)

Marked by CJL

**Marks**

(a)



The point  $P(\sec \theta, \tan \theta)$  lies on the hyperbola  $H$  with equation  $x^2 - y^2 = 1$ . A vertical line through  $P$  intersects with an asymptote at  $S$  and with the  $x$ -axis at  $T$  as shown. A normal to  $H$  at  $P$  intersects the  $x$ -axis at  $R$ . The point  $F$  is a foci of  $H$ .


- (i) Show that the equation of the normal to  $H$  at the point  $P$  is  $y = -\sin \theta x + 2 \tan \theta$ . 2
- (ii) Show that  $RS = \sqrt{2}RT$ . 3
- (iii) Find the coordinates of the point  $U$  which lies on  $SR$  such that  $TU$  is parallel to the asymptote on which  $S$  lies. 2
- (iv) For what values of  $\theta$  will  $FU$  be the perpendicular bisector of  $SR$ ? 2
- (b) Let  $\omega = \cos \frac{\pi}{5} + i \sin \frac{\pi}{5}$ .
- (i) Write down in terms of  $\omega$ , and the positive integer  $k$ , all the solutions of the equation  $z^{10} - 1 = 0$ . 2
- (ii) Prove that  $\omega + \omega^2 + \omega^3 + \dots + \omega^{10} = 0$ . 2
- (iii) The quadratic equation  $x^2 + bx + c = 0$ , where  $b$  and  $c$  are real, has the root  $\omega + \omega^4$ . Find the other root in terms of  $\omega$ . 2

**Question 7** (15 marks)

Marked by CJL

**Marks**

(a)



In the diagram,  $\ell$  is a circle with exterior point  $P$ . Tangents from  $P$  are drawn to meet  $\ell$  at points  $A$  and  $E$ . The point  $C$  is the centre of  $\ell$ . The line  $BP$  passes through  $C$ . The line  $AD$  passes through  $B$ . The line  $CF$  passes through  $E$ .  $AP$  is parallel to  $DF$ .

(i) Show that  $ACEP$  is a cyclic quadrilateral. 1

(ii) Use a double angle formula to show that  $DE = \frac{DF(EP^2 - CE^2)}{CP^2}$  2

(iii) Use the sine rule to show that  $\frac{AB}{BD} = \frac{AP}{DP}$  2



- |     |  | <b>Marks</b> |
|-----|--|--------------|
| (b) | (i) Draw the graph of $y = \ln(x + 1)$   | <b>1</b>     |
|     | (ii) Hence explain why   | <b>1</b>     |
|     | $\int_0^n \ln(x + 1) dx < n \ln(n + 1), \quad n = 1, 2, 3, \dots$  |              |
|     | (iii) Use integration by parts to show that  | <b>3</b>     |
|     | $\int_0^n \ln(x + 1) dx = \ln(n + 1)^{n+1} - n$  |              |
|     | (iv) Hence deduce that $\ln(n + 1) < n$  | <b>1</b>     |
|     | (v) Show that $\sum_{k=1}^n \frac{1}{2} \ln(k + 1) = \frac{1}{2} \ln(n + 1)!$  | <b>1</b>     |
|     | (vi) Use the results from parts (iii) and (v) together with your graph to deduce that $n! < \left(\frac{n + 1}{e}\right)^{2n} (n + 1)$ | <b>3</b>     |

**Question 8** (15 marks)

**Marked by SKB**

**Marks**

- (a) (i) For all real, positive numbers  $a$  and  $b$ , where  $a > b$  show that **4**
- ( $\alpha$ )  $a + b > 2\sqrt{ab}$
- ( $\beta$ )  $b^2 - a^2 < 2\sqrt{ab}(b - a)$
- (ii) Hence deduce that  $a > c$  given that  $c$  is a positive real number and **4**
- $$\sqrt{a}(b - a) + \sqrt{c}(c - b) > \frac{c^2 - a^2}{2\sqrt{b}}$$
- (b) If  $h(n) = n^4 + 6n^2 + 9$
- (i) show that  $h(n + 2) - h(n) = 8(n + 1)(n^2 + 2n + 5)$  **3**
- (ii) hence prove by mathematical induction that  $h(n)$  is divisible by 8 if  $n$  is an odd positive integer. **4**

**END OF EXAM**

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

**NOTE :**  $\ln x = \log_e x, \quad x > 0$