



TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION 2003

MATHEMATICS

EXTENSION II

*Time Allowed – 3 Hours
(Plus 5 minutes reading time)*

All questions may be attempted

All questions are of equal value

In every question, show all necessary working

Marks may not be awarded for careless or badly arranged work

**Standard integral tables are provided for your convenience.
Approved silent calculators may be used.**

**The answers to all questions are to be returned in separate bundles
clearly labelled Question 1, Question 2, etc. Each bundle must show your
candidate number.**

JRAHS TRIAL - EXT II 2003

Question 1:

- (a) The complex numbers $z_1 = \frac{a}{1+i}$ and $z_2 = \frac{b}{1+2i}$ where a and b are real, satisfy the condition $z_1 + z_2 = 1$. Find the value of a and b . 3
- (b) The complex number z has modulus r and argument θ where $0 \leq \theta \leq \pi$. Find in terms of r and θ the modulus and arguments of
- (i) z^2 1
- (ii) $\frac{1}{z}$ 1
- (iii) iz 1
- (c) (i) Sketch (without using calculus) the curve $y = \frac{x^2 + 2x - 3}{x - 2}$ clearly showing its intercepts with the coordinate axes and the position of all its asymptotes. 5
- (ii) Find the area bounded by the curve $y = \frac{x^2 + 2x - 3}{x - 2}$ and the x -axis. 4

Question 2: (START A NEW PAGE)

- (a) Evaluate:
- (i) $\int_0^{\frac{\pi}{2}} \cos \theta \sin^3 \theta \, d\theta$. 2
- (ii) $\int_0^3 \frac{\sqrt{x}}{1+x} \, dx$. (Let $u^2 = x$). 3
- (iii) $\int_0^{\frac{\pi}{2}} \frac{1}{5+3\cos\theta} \, d\theta$ 4
- (b) Given that $I_n = \int_0^{\frac{\pi}{2}} x^n \sin x \, dx$.
- (i) Prove that $I_n + n(n-1)I_{n-2} = n\left(\frac{\pi}{2}\right)^{n-1}$. 4
- (ii) Evaluate $\int_0^{\frac{\pi}{2}} x^4 \sin x \, dx$. 2

Question 3: (START A NEW PAGE)

- (a) Sketch the ellipse $9x^2 + 25y^2 = 225$ clearly showing: 4
- (i) the coordinates of the intercepts with the x and y -axes,
 - (ii) the coordinates of the foci,
 - (iii) the equation of the directrices.
- (b) Prove that the curves $x^2 - y^2 = c^2$ and $xy = c^2$ meet at right angles. 4
- (c) The point $P(a \cos \theta, b \sin \theta)$ lies on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the ellipse meets the x -axis at the points A and A' .
- (i) Prove that the tangent at P has the equation $\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$. 3
 - (ii) The tangent at P meets the tangents from A and A' at points Q and Q' respectively. Find the coordinates of Q and Q' . 2
 - (iii) Prove that the product $AQ \times A'Q'$ is independent of the position of P . 2

Question 4: (START A NEW PAGE)


- (a) Prove that $\frac{d}{dx} \left[\sqrt{bx - x^2} + \frac{b}{2} \cos^{-1} \left(\frac{2x - b}{b} \right) \right] = -\sqrt{\frac{x}{b - x}}$ for $x \geq 0$. 3
- (b) A particle of mass m is attracted towards the origin by a force of magnitude $\frac{\mu m}{x^2}$ for $x \neq 0$, where the distance from the origin is x and μ is a positive constant.
- (i) If the particle starts from rest at a distance b to the right of the origin, show that its velocity v is given by $v^2 = 2\mu \left(\frac{b - x}{bx} \right)$. 3
 - (ii) Find the time required for the particle to reach a point halfway towards the origin. 4
- (c) Using the Principle of mathematical induction, prove that $(x + 1)^n - nx - 1$ is divisible by x^2 for all integer $n \geq 2$. 5

Question 5: (START A NEW PAGE)

- (a) (i) Using the substitution $x = 2 \sin \theta$, prove that $\int_0^2 (4-x^2)^{\frac{3}{2}} dx = 16 \int_0^{\frac{\pi}{2}} \cos^4 \theta d\theta$ 2

- (ii) A solid (see diagram) sits on a semi-circular base of radius 2 units. Vertical cross-sections perpendicular to the diameter of the semi-circle are right-angled triangles with their heights being bounded by the parabola $y = 4 - x^2$. By slicing the solid perpendicular to the x -axis, show that the volume (V unit³) of the solid formed is given by

$$V = \int_0^2 (4-x^2)^{\frac{3}{2}} dx$$



- (iii) Find the volume of the solid. 5

- (b) A tourist is walking along a straight road. At one point he observes a vertical tower standing on a large flat plain. The tower is on a bearing 053° with an angle of elevation of 21° . After walking 230 metres, the tower is on a bearing 342° with an angle of elevation of 26° .

- (i) Draw a neat diagram showing the above information. 1
- (ii) Find the height of the tower correct to the nearest metre. 5

Question 6: (START A NEW PAGE)

- (a) The tangent to the hyperbola $xy = c^2$ at the point $T\left(ct, \frac{c}{t}\right)$ meets the x and y axes at F and G respectively and the normal at T meets the line $y = x$ at H .

- (i) Show that the tangent at T is $x + t^2y = 2ct$. 3

- (ii) Show that the normal at T is $t^3x - ty = c(t^4 - 1)$. 2

- (iii) Prove that $FH \perp HG$. 6

- (b) The area bounded by the curve $y = \frac{\ln x}{\sqrt{x}}$ and the x -axis for $1 \leq x \leq e$ is rotated through one revolution about the y -axis. Using the method of cylindrical shells find the volume of the solid. 4


Question 7: (START A NEW PAGE)

(a) In a state swimming championships, 12 swimmers (including the Jones twins) are chosen to represent their club and are divided into three teams of four swimmers to form 3 relay teams. Find the number of ways this can be done:

- (i) with no restrictions. 2
- (ii) if the Jones twins (Angela and Bethany) are not to be in the same relay team. 3

(b) The ends of a light string are fixed at 2 points A and B with B directly below A, as shown in the diagram. The string passes through a small ball of mass m which is then fastened to the string at point P. The angle PAB is θ and the distance from P to AB is r .

Suppose now that the ball revolves in a horizontal circle about the vertical through AB with constant angular velocity ω and while this happens both sections (AP and BP) of the string are taut and the angle APB is a right angle.




- (i) Draw a diagram showing the forces acting on the ball. 2
- (ii) Show that the tensions T_1 and T_2 in the sections of the string AP and BP respectively are $T_1 = m(r\omega^2 \sin \theta + g \cos \theta)$ and $T_2 = m(r\omega^2 \cos \theta - g \sin \theta)$. 4
- (iii) Given that $AB = 100\text{cm}$ and $AP = 80\text{cm}$, show that $\omega^2 > \frac{25g}{16}$. 2
- (iv) Suppose that the ball is free to slide on the string. Show that the condition for the ball to remain at point P on the string is $\omega^2 = \frac{175g}{12}$. 2

Question 8: (START A NEW PAGE)

- (a) (i) If $t = \tan x$ prove that $\tan 4x = \frac{4t(1-t^2)}{t^4 - 6t^2 + 1}$ 2
- (ii) If $\tan x \tan 4x = 1$ deduce that $5t^4 - 10t^2 + 1 = 0$. 1
- (iii) Prove that $x = 18^\circ$ and $x = 54^\circ$ satisfy the equation $\tan x \tan 4x = 1$. 2
- (iv) Deduce that $\tan 54^\circ = \sqrt{\frac{5+2\sqrt{5}}{5}}$. 3

- (b) ABCD is a cyclic quadrilateral and E is on BD such that $\angle DAE = \angle BAC$.



- (i) Copy the diagram onto your answer sheet and prove that $\triangle ABE$ and $\triangle ADC$ are similar. 2
- (ii) Prove that $AB \times CD = AC \times BE$. 1
- (iii) Hence by proving that another pair of triangles are similar, deduce that $AB \times CD + AD \times BC = AC \times BD$. 4

THE END