


JRAHS Trial HSC 2006 Extension 2

Question 1 (15 Marks)	Marks
(a) Find $\int \frac{2x}{1+2x} dx$.	2
(b) How many different ways are there of choosing a cricket 11 from a team of 15 players, if there can only be at most, one of the 2 Lee brothers and 2 of 3 Abey brothers?	3
(c) Find $\lim_{x \rightarrow -5} \frac{\sqrt{20-x} - 5}{5+x}$.	2
(d) Find the sum of all the coefficients of powers of z in the expansion $(1+z)^8$.	2
(e) (i) Find the equation of the tangent to the hyperbola $3x^2 - y^2 = 12$ at the point $T(x_1, y_1)$.	3
(ii) This tangent meets the line $d(x=1)$ in the point M . Prove that $FM \perp FT$, where F is the focus $(4, 0)$.	3

Question 2 (15 Marks) START A NEW PAGE

(a) (i) Express $z = \sqrt{2} - i\sqrt{2}$ in modulus – argument form.	2
(ii) Hence, write z^{22} in the form of $a + ib$, where a and b are real.	3
(b) If a, b, c are real and unequal and that $a^2 + b^2 > 2ab$ deduce that	
(i) $a^2 + b^2 + c^2 > ab + bc + ca$	2
(ii) If $a + b + c = 6$ show that $ab + bc + ca < 12$	2
(c) Let $I_n = \frac{1}{n!} \int_0^1 x^n e^{-x} dx$ for $n \geq 0$.	
(i) Use integration by parts to show that $I_n = I_{n-1} - \frac{e^{-1}}{n!}$	3
(ii) Hence, evaluate I_4 .	3

(a) The diagram below is a sketch of the function $y = f(x)$.



On separate diagrams sketch neatly:

- (i) $y = |f(x)|$ 2
- (ii) $y = e^{f(x)}$ 2
- (iii) $y = \ln(f(x))$ 2
- (b) Reduce the polynomial $x^4 - 2x^2 - 15$ over the rational and the complex field. 2
- (c) If z_1, z_2 are 2 complex numbers such that $|z_1 + z_2| = |z_1 - z_2|$. 4
 With the aid of a diagram, show that $\arg(z_1)$ and $\arg(z_2)$ differ by either $\frac{\pi}{2}$ or $\frac{3\pi}{2}$.
- (d) Evaluate $\int_0^{\frac{\pi}{2}} \frac{1 - \tan x}{1 + \tan x} dx$. 3

Question 4 (15 Marks)

START A NEW PAGE


Marks

- (a) A string 50 cm of length can just sustain a weight of mass 20 kg without breaking. A mass of 4 kg is attached to one end of the string and revolves uniformly on a smooth horizontal table; the other end is being fixed to a point on the table. **3**

Find the greatest number of complete revolutions the mass can make in a minute without breaking the string. [Use acceleration due to gravity as 9.8 m/s^2]

- (b) Evaluate $\int_a^{a^2} \frac{dx}{x \ln x}$, leaving your answer in simplified, exact form. **3**

- (c)



The area between the graphs of $y = \cos^2 x$ and $y = \cos 2x$ for $0 \leq x \leq \frac{\pi}{2}$ **6**

and $x = \frac{\pi}{2}$ is rotated about the y – axis. By considering cylindrical shells, find volume of revolution of the solid formed, in terms of π .

- (d) Use the properties of odd and even functions to evaluate $\int_{-4}^4 \cos x(e^x - e^{-x}) dx$. **3**

Question 5 (15 Marks) START A NEW PAGE

(a) (i) Find A and B for which $\frac{x^4}{x^2+1} = A(x^2-1) + \frac{B}{x^2+1}$. 2

(ii) Show that this result may be used in the integration of the function $x^3 \tan^{-1} x$. 5

Hence, evaluate the integral $\int_0^1 x^3 \tan^{-1} x \, dx$.

(b) A particle P is projected vertically upwards from the surface of the Earth with initial speed u . The acceleration due to gravity at any point on its path is inversely proportional to the square of its distance from the centre of the Earth.




Diagram not to scale

(i) Prove that the speed v in any position x is given by $v^2 = u^2 - 2gR + \frac{2gR^2}{x}$, 4

where R is the radius of the Earth and g is the acceleration due to gravity at the surface of the Earth.


(ii) If $u = \sqrt{2gR}$, find the time taken to reach a height $3R$ above the surface of the earth, in terms of g and R . 4

Question 6 (15 Marks)

START A NEW PAGE

Marks

- (a) A total of five players is selected at random from four sporting teams. Each of the teams consists of 10 players numbered from 1 to 10. What is the probability that the five selected players contain at least four players from the same team? Clearly explain your answer. 4
- (b) A particle P of mass m is attached by two equal, light inextensible strings of length l , to two points Q and S , distant h units apart in the same vertical line; S is directly below Q . P rotates in a horizontal circle with uniform angular velocity, ω .



- (i) Prove that the tension in the string PQ is $ml\left(\frac{1}{2}\omega^2 + \frac{g}{h}\right)$ where g is the acceleration due to gravity. 5
- (ii) Find the tension in the string PS . 1
- (iii) Show that for both strings to remain stretched then $\omega > \sqrt{\frac{2g}{h}}$. 2
- (iv) If the tensions in the string are in the ration of 2:1, find the period of the motion of the particle. 3

Question 7 (15 Marks)

START A NEW PAGE


Marks

- (a) α, β, γ are non – zero roots of the equation $x^3 + px + q = 0$.
Find the equation whose roots are

(i) $\alpha^3, \beta^3, \gamma^3$ 3

(ii) $\frac{\alpha}{\beta\gamma}, \frac{\beta}{\alpha\gamma}, \frac{\gamma}{\alpha\beta}$ 4

- (b) The diagram below shows the hyperbola $xy = c^2$. The point $P\left(ct, \frac{c}{t}\right)$ lies on the curve, where $t \neq 0$. The normal at P intersects the straight line $y = x$ at N . O is the origin.



- (i) Prove that the equation of the normal at P is 1

$$y = t^2x + \frac{c}{t} - c^3.$$

- (ii) Find the coordinates of N . 1

- (iii) Show that $\triangle OPN$ is isosceles. 2


- (c) (i) Draw a neat sketch of the graph of $y = \frac{4}{x} - x$. 2

- (ii) Draw a neat sketch of the curve $y = \sqrt{f(x)}$. 2

Question 8 (15 Marks)**START A NEW PAGE****Marks**

- (a) Ben's Hot Wheel travels along an inclined ramp to jump over a truck. The Hot Wheel travels at an initial speed V m/s inclined at an angle θ to the horizontal at O , and acceleration due to gravity is g m/s².

The Wheel's trajectory is given by $y = x \tan \theta - \frac{gx^2 \sec^2 \theta}{2V^2}$.



- (i) Write this equation in the general form of a parabola: $(x - h)^2 = 4a(y - k)$, where a, h, k are constants. **3**
- (ii) Calculate the angle of projection, θ , if the range is three times the width of the truck and the top of the truck passes through the focus of equation in part (a). **3**
- (b) Given p and q are positive real numbers,
- (i) Prove that: $\frac{1}{2}(p + q) \geq \sqrt{pq}$. **1**
- (ii) Hence, deduce that: $\sqrt{p} \leq \frac{1}{2} \left(\frac{p}{\sqrt{q}} + \sqrt{q} \right)$. **1**

Question 8 Part (c) continued on the next pag

(c) Given $a_1, a_2, a_3, \dots, a_n$ and $b_1, b_2, b_3, \dots, b_n$ are positive real numbers, where $A_n = a_1 + a_2 + a_3 + \dots + a_n$ and $B_n = b_1 + b_2 + b_3 + \dots + b_n$, are such that $a_1, a_2, \dots, a_n > 0$, $b_1, b_2, \dots, b_n > 0$ and $A_r \leq B_r$, for $r = 1, 2, 3, \dots, n$.

(i) Prove, by mathematical induction for $n = 1, 2, 3, \dots$, that: **3**

$$\begin{aligned} & \frac{1}{\sqrt{b_n}} B_n + \left(\frac{1}{\sqrt{b_{n-1}}} - \frac{1}{\sqrt{b_n}} \right) B_{n-1} + \left(\frac{1}{\sqrt{b_{n-2}}} - \frac{1}{\sqrt{b_{n-1}}} \right) B_{n-2} + \dots + \left(\frac{1}{\sqrt{b_1}} - \frac{1}{\sqrt{b_2}} \right) B_1 \\ & = \sqrt{b_1} + \sqrt{b_2} + \sqrt{b_3} + \dots + \sqrt{b_n}. \end{aligned}$$

(ii) Hence, given: $\frac{a_1}{\sqrt{b_1}} + \frac{a_2}{\sqrt{b_2}} + \frac{a_3}{\sqrt{b_3}} + \dots + \frac{a_n}{\sqrt{b_n}} =$

$$\frac{1}{\sqrt{b_n}} A_n + \left(\frac{1}{\sqrt{b_{n-1}}} - \frac{1}{\sqrt{b_n}} \right) A_{n-1} + \left(\frac{1}{\sqrt{b_{n-2}}} - \frac{1}{\sqrt{b_{n-1}}} \right) A_{n-2} + \dots + \left(\frac{1}{\sqrt{b_1}} - \frac{1}{\sqrt{b_2}} \right) A_1,$$

show that: $\sum_{r=1}^n \frac{a_r}{\sqrt{b_r}} \leq \sum_{r=1}^n \sqrt{b_r}$. **1**

(iii) Deduce that: $\sum_{r=1}^n \sqrt{a_r} \leq \sum_{r=1}^n \sqrt{b_r}$. **3**