

THE KING'S SCHOOL

---

2003  
Higher School Certificate  
Trial Examination

**Mathematics Extension 2**

**General Instructions**

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

**Total marks – 120**

- Attempt Questions 1-8
- All questions are of equal value

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

**Marks**

**Question 1 (15 marks)** Use a SEPARATE writing booklet.

(a) Find

(i)  $\int \frac{1+x+x^2}{1+x^2} dx$

2

(ii)  $\int \frac{x^2}{1+x^2} dx$

2

(b) Use integration by parts to evaluate

$$\int_0^1 2x \tan^{-1} x \, dx$$

3

(c) Find  $\int_0^1 \frac{x-3}{(x^2+1)(3x+1)} dx$ , giving your answer in simplest exact form.

4

(d)  $u_n = \int_0^1 \frac{x^n}{1+x^2} dx, n \geq 0$

(i) Show that  $u_{n+2} + u_n = \frac{1}{n+1}$

2

(ii) Hence, evaluate  $\int_0^1 \frac{x^3}{1+x^2} dx$

2

**End of Question 1**

**Marks**

**Question 2 (15 marks)** Use a SEPARATE writing booklet.

(a)  $u = 2 + ai, v = a + 2i$ , where  $a$  is a real number.

Find in the form  $x + iy$ ,

(i)  $uv$

2

(ii)  $(uv)^{-1}$

1

(b) (i) Express  $z = -2\sqrt{3} + 2i$  in modulus-argument form

2

(ii) Hence, find  $z^3$  in the form  $x + iy$

2

(c) Sketch the region in the complex plane where

$$|z-i| \leq |z+1|$$

3

(d) Consider the equation  $(a+ib)^2 = 1+2i$ ,  $a, b$  real

(i) Show that  $a^2 + b^2 = \sqrt{1^2 + 2^2}$

1

(ii) Hence, or otherwise, find the value of  $a^2$


2

**Question 2 continues on next page**

Question 2 (continued)

Marks

- (e) In the complex plane, A is the point (1,0) and the complex number  $\overline{AB}$  is  $w-1$ .  $\triangle ABC$  is isosceles and right-angled at B. O is the origin.



Find, in terms of  $w$ , the complex numbers

- (i)  $\overline{CB}$   
 (ii)  $\overline{OC}$

1

1

End of Question 2

Marks

Question 3 (15 marks) Use a SEPARATE writing booklet.

- (a) (i) Sketch on the same axes the graphs of

$$y = |x-1| \text{ and } y = 2x-x^2$$

2

- (ii) Use (i) to show on separate diagrams, the graphs of

( $\alpha$ )  $y = \frac{|x-1|}{2x-x^2}$ , showing any asymptotes

3

( $\beta$ )  $y = \frac{2x-x^2}{|x-1|}$ , showing any asymptotes

3

- (b) Consider the function  $f(x) = \tan^{-1} x - \frac{x}{1+x^2}$

- (i) Show that  $f$  is an odd function.

1

- (ii) Find  $f'(x)$

2

- (iii) Show that  $f(x) > 0$  if  $x > 0$

2

- (iv) Sketch the graph of  $y = \tan^{-1} x - \frac{x}{1+x^2}$


2

End of Question 3

Question 4 (15 marks) Use a SEPARATE writing booklet.

- (a) Find the gradient of the tangent to the curve  $x^3 + y^2 + xy = 0$  at the point  $(-2, 4)$  3
- (b)  $P(x_1, y_1)$  is a point in the first quadrant on the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a > b > 0$

$S$  and  $S'$  are the foci of the ellipse.  $O$  is the origin.



- (i) Show that the equation of the normal at  $P(x_1, y_1)$  is  $a^2 y_1(x - x_1) = b^2 x_1(y - y_1)$  2
- (ii) The normal at  $P$  meets the major axis at  $N$ .  
Prove that the  $x$  coordinate of  $N$  is  $e^2 x_1$ , where  $e$  is the eccentricity of the ellipse. 2
- (iii) Deduce that  $N$  lies between  $O$  and  $S$ . 2
- (iv) Show that  $NS = eSP$  and  $NS' = eS'P$  3
- (v) Using the sine rule in  $\triangle PSN$  and  $\triangle PS'N$ , or otherwise, prove that  $PN$  bisects  $\angle SPS'$  3

End of Question 4

Question 5 (15 marks) Use a SEPARATE writing booklet.

- (a) Four married couples are to be seated at a circular table.
- (i) How many arrangements are possible if the men and women are to be separated? 2

- (ii) For the arrangements in (i), find the probability that no woman is sitting next to her husband. 2

- (b) The equation  $x^3 + ax^2 + bx + c = 0$  has one root the sum of the other two roots.  
Prove that  $a^3 - 4ab + 8c = 0$  4

- (c) (i) By considering the circle  $x^2 + y^2 = a^2$ , or otherwise, find

$$\int_0^a \sqrt{a^2 - x^2} \, dx$$

2

- (ii) The ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a > b > 0$ , is revolved about the line  $y = a$ .

By considering slices perpendicular to the line  $y = a$ , find the volume of the solid of revolution generated.

5

End of Question 5

**Question 6 (15 marks)** Use a SEPARATE writing booklet.

- (a) A particle of mass  $m$  kg falls vertically from rest from point O in a medium whose resistance is  $mkv$ , where  $k$  is a positive constant and  $v$  is its velocity in m/s. After  $t$  seconds the particle has fallen  $x$  metres.  
 $g$  m/s<sup>2</sup> is the acceleration due to gravity.

(i) Show that  $\frac{dv}{dt} = g - kv$

- (ii) Find the terminal velocity,  $V$  m/s, of the particle.

(iii) Use integration to prove that  $v = \frac{g}{k} (1 - e^{-kt})$

- (iv) Find the distance the particle has fallen when its velocity is one half of its terminal velocity.

(b)  $\alpha, \beta$  are the two complex roots of the equation  $x^3 + 5x + 1 = 0$

- (i) Explain why  $\alpha, \beta$  are complex conjugates.

(ii) Show that the real root is  $\frac{-1}{|\alpha|^2}$

(iii) Show that  $\alpha\beta$  is a root of the equation  $x^3 - 5x^2 - 1 = 0$

End of Question 6

Marks

1

1

3

4

1

2

3

Marks

**Question 7 (15 marks)** Use a SEPARATE writing booklet.

- (a) By mathematical induction it is easy to show that

$$1^2 - 2^2 + 3^2 - \dots - (2n)^2 = -n(2n+1)$$

If, further, it is known that

$$1^2 + 2^2 + 3^2 + \dots + (2n)^2 = \frac{n}{3}(2n+1)(4n+1),$$

deduce that

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n}{6}(n+1)(2n+1)$$

(Do not use induction)

3

- (b) (i) Using the substitution  $t = \tan \frac{x}{2}$ , or otherwise, evaluate

$$\int_0^{\frac{\pi}{2}} \frac{dx}{1 + \sin x}$$

3

- (ii) Let  $F(x)$  be a primitive function of  $f(x)$ .

Using this, or otherwise, show that

$$\int_0^{2a} f(x) dx = \int_0^a f(x) + f(2a-x) dx$$

2

- (iii) Deduce  $\int_0^{\frac{\pi}{2}} \frac{x}{1 + \sin x} dx$

3

Question 7 continues next page

Question 8 (15 marks) Use a SEPARATE writing booklet.

(a) The roots of  $z^n = 1$ ,  $n$  a positive integer, are

$$z_k = \cos \frac{2\pi k}{n} + i \sin \frac{2\pi k}{n}, \quad k = 1, 2, \dots, n$$

(i) Show that  $z_k^n = z_1^{kp}$ ,  $p$  a positive integer

(ii) If  $z_k$  is such that  $z_k, z_k^2, z_k^3, \dots, z_k^n$  generates all the roots of  $z^n = 1$ , then  $z_k$  is called a primitive root of  $z^n = 1$

( $\alpha$ ) Show that  $z_1$  is a primitive root of  $z^n = 1$

( $\beta$ ) Show that  $z_5$  is a primitive root of  $z^6 = 1$

( $\gamma$ ) Suppose the highest common factor of  $n$  and  $k$  is  $h$ , i.e.  $n = ph$  and  $k = qh$ ,  $p, q$  integers.

Show that for  $z_k$  to be a primitive root of  $z^n = 1$ , then  $h = 1$

(b) (i) Show that  $\sum_{k=0}^{n-1} (1-x)^k = \frac{1-(1-x)^n}{x}$ ,  $x \neq 0$


(ii) Deduce that  $\sum_{k=0}^{n-1} (1-x)^k = \sum_{k=1}^n (-1)^{k-1} \binom{n}{k} x^{k-1}$

(iii) Explain or show why  $\int \sum_{k=0}^{n-1} (1-x)^k dx = \sum_{k=0}^{n-1} \int (1-x)^k dx$

(iv) Deduce that  $\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{1}{k} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$

Question 7 (continued)

(c) A mass  $m$  at  $P$  is freely joined to two equal light rods  $PQ$  and  $PR$  of length  $a$ . The end  $Q$  of  $PQ$  is pivoted to a fixed point  $Q$  and the end  $R$  of  $PR$  is freely joined to a ring of mass  $m$  which slides on a smooth vertical pole. If  $P$  rotates in a horizontal circle with uniform angular velocity  $\omega$ , show the angle of inclination of the rods  $PQ$  and  $PR$  to the vertical is  $\tan^{-1} \left( \frac{r\omega^2}{3g} \right)$ .  $T_1, T_2$  are tensions in the rods,  $N$  is the normal reaction of  $QR$  on the ring  $R$ .



End of Question 7

2

1

2

2

2

2

1

3

4