

Trial Use

Pyndle LC. 2004 Trial Ext 2

Question 1. Use a SEPARATE Writing Booklet

Marks

(a) Find $\int 3x \sec^2(x^2) dx$

[2]

(b) Find $\int \frac{e^{-x}}{\sqrt{1-e^{-x}}} dx$

[2]

(c) Find $\int \frac{\tan^2 x}{\sin^2 x} dx$

[2]

(d) Find $\int \cos^{-1} \theta d\theta$

[3]

(e) Let $I_n = \int_0^{\pi} x^n \sin x dx$ where n is an integer

(i) Show that $I_n = \pi^n - n(n-1)I_{n-1}$ for $n \geq 2$

[3]

(ii) Hence evaluate I_5

[3]

Question 2 Use a SEPARATE Writing Booklet

Marks

(a) Given that $W = \frac{3+i}{1-2i}$, express the following in the form $a+ib$

(i) $|W|$

[2]

(ii) \bar{W}

[1]

(iii) W^{-1}

[1]

(b) Illustrate with a diagram and describe in Geometric terms the locus, represented by the following:

(i) $|-z| = 3$

[2]

(ii) $\frac{\bar{z}z}{2} = \bar{z} + z$

[2]

(c) Let z be a complex number such that $\arg(z) = \theta$, where $\frac{\pi}{2} < \theta < \pi$, and $|z| = 1$

Sketch z^2 and z on an Argand Diagram and find in terms of θ the values of

(i) $\frac{2}{|1+z^2|}$

[3]


(ii) $\arg\left(\frac{2}{1+z^2}\right)$

[4]

Question 3. Use a SEPARATE Writing Booklet

Marks

- (a) On your answer page make a rough copy of the sketch of $f(x)$ below. Label the turning points A, B, C and label the cusp D as indicated. On the same set of axes sketch the derivative function, $f'(x)$ [3]



- (b) Sketch the curves on two separate diagrams, show the equations of any asymptotes, and show any intercepts on the axes.

(i) $y = \frac{5-x}{x}$ [2]

(ii) $y = \frac{25-x^2}{x^2}$ [2]

Hence or otherwise sketch on another two separate diagrams, the curves

(iii) $y = \left| \frac{5-x}{x} \right|$ [2]


(iv) $y^2 = \frac{25-x^2}{x^2}$ [3]

- (c) Sketch and label $y = \sin^2(2x)$ for $0 \leq x \leq \pi$ and on the same axes sketch and label $y = \ln(\sin^2(2x))$ for $0 \leq x \leq \pi$. [3]

Question 4. Use a SEPARATE Writing Booklet

Marks

- (a) A large shiny metal teapot appears to be circular when viewed from above. The same teapot appears to be elliptical when viewed from the side (as shown). [6]




- Use the method of shells to find the volume of this shiny metal teapot and then express its capacity in litres correct to 2 decimal places.

- (b) With the aid of a sketch and careful integration show that the area enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is πab square units. [3]

Question 4 continues next page


Question 4. continued

- (c) The horizontal cross-section of a vase is a regular octagon. The maximum width $2x$ of the octagonal cross-section is 10cm at the base and 20cm at the top.



The horizontal cross-section is shown opposite

The vertical cross-section is shown opposite.




- (i) Find an expression for the area of the horizontal cross-section when the maximum width is $2x$ [2]
 (ii) Find the volume of the vase using the method of parallel cross-sections. [leave your answer in cm^3] [4]

Question 5. Use a SEPARATE Writing Booklet

Marks

- (a) A square-based pyramid, of base length, s and perpendicular height, h is inscribed in a sphere of radius r .
 (i) Show that $s^3 = 4hr - 2h^2$ [2]
 (ii) Find h in terms of r so that the pyramid has maximum volume. [3]
 (iii) Find an expression for the maximum volume in terms of r . [2]

- (b) The diagram below represents a Cyclic Quadrilateral $ABCD$.



Bisectors of the angles have been drawn, forming a smaller quadrilateral $EFGH$. Copy the diagram onto your answer page and Prove that $EFGH$ is also a Cyclic Quadrilateral [5]

- (c) Solve for x [3]
 $\tan^{-1} 5x - \tan^{-1} 3x = \tan^{-1} \frac{1}{4}$

Question 6.	Use a SEPARATE Writing Booklet	Marks
(a)	(i) Express the complex cube roots of unity, ω and ω^2 in the form $re^{i\theta}$	[1]
	(ii) Using Argand diagram or otherwise, show that $\omega + \omega^2 + \omega^3 = 0$	[3]
	(iii) Prove that if $P(x) = p_0 + p_1x + p_2x^2 + \dots + p_nx^n$ has a root of multiplicity m , then $P'(x)$ has a root of multiplicity $(m-1)$.	[3]
	(iv) Prove that ω , a complex cube root of unity is a repeated root of the polynomial $P(x) = 5x^3 + 7x^4 + 9x^5 + x^2 - x - 3$	[4]
(b)	If α, β, γ are the roots of $x^3 - 9x + 9 = 0$, show that $(\alpha - 1)(\beta - 1)(\gamma - 1) = -1$	[4]

Question 7.	Use a SEPARATE Writing Booklet	Marks
(a)	(i) Write down the parametric equations that correspond to the Cartesian equation $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	[1]
	(ii) Show that the greatest area of a rectangle inscribed in an ellipse is $2ab$	[2]
(b)	For the hyperbola $\frac{x^2}{16} - \frac{y^2}{9} = 1$ find	
	(i) Eccentricity	[1]
	(ii) Coordinates of the foci	[1]
	(iii) Equations of the directrices	[1]
	(iv) Equation of the tangent to the hyperbola at the point $(5, -2\frac{1}{4})$	[2]
	(v) Show foci, directrices, asymptotes and the tangent on a neat sketch.	[4]
(c)	An hyperbola has asymptotes $y = \pm x$ and it passes through the point $(-3, -2)$.	
	(i) Find the equation of the hyperbola	[1]
	(ii) Find the length of the transverse axis	[1]
	(iii) Explain why this hyperbola is rectangular	[1]

Question 8. Use a SEPARATE Writing Booklet

Marks

- (a) The positive integers are bracketed as follows:
(1), (2,3), (4,5,6),
where there are r integers in the r^{th} bracket.
Prove that the sum of the integers in the r^{th} bracket is $\frac{1}{2}r(r^2 + 1)$. [5]
- (b) If $a > 0, b > 0, c > 0$, show that $a^2 + b^2 + c^2 \geq ab + bc + ca$
and state the condition of equality. [3]
- (c) (i) Express $1 + x + x^2 + x^3 + x^4 + x^5$ as a product of real factors [1]
(ii) Prove that the equation $c + x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \frac{x^5}{5} + \frac{x^6}{6} = 0$
has no real roots if $c > \frac{37}{60}$ [6]

END OF PAPER