

St George Girls High School

Trial Higher School Certificate Examination

2012



# Mathematics Extension 1

## General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using blue or black pen.
- Write your student number on each booklet.
- Board-approved calculators may be used.
- The mark allocated for each question is listed at the side of the question.
- Marks may be deducted for careless or poorly presented work.

## Total Marks – 100

### Section I – Pages 2 – 4 10 marks

- Attempt Questions 1 – 10.
- Allow about 15 minutes for this section.
- Answer on the sheet provided.

### Section II – Pages 5 – 12 60 marks

- Attempt Questions 11 – 14
- Allow about 2 hours 45 minutes for this section.
- Begin each question in a new booklet.
- Show all necessary working in Questions 11 – 14.

Students are advised that this is a Trial Examination only and does not necessarily reflect the content or format of the Higher School Certificate Examination.

Section I – (10 marks)

Marks

Answer this section on the answer sheet provided at the back of this paper.

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

1. The value of  $\lim_{x \rightarrow 0} \frac{\sin 4x}{9x}$  is:

A.  $2\frac{1}{4}$

B. 1

C.  $\frac{4}{9}$

D. 0

2. For the function  $f(x) = 3 \sin^{-1}\left(\frac{x}{4}\right)$  the domain and range of  $y = f(x)$  are:

A. domain  $\left\{x: -\frac{3\pi}{2} \leq x \leq \frac{3\pi}{2}, x \in \mathbb{R}\right\}$

range  $\{y: -4 \leq y \leq 4, y \in \mathbb{R}\}$

B. domain  $\{x: -1 \leq x \leq 1, x \in \mathbb{R}\}$

range  $\{y: -3 \leq y \leq 3, y \in \mathbb{R}\}$

C. domain  $\{x: -3 \leq x \leq 3, x \in \mathbb{R}\}$

range  $\left\{y: -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}, y \in \mathbb{R}\right\}$

D. domain  $\{x: -4 \leq x \leq 4, x \in \mathbb{R}\}$

range  $\left\{y: -\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}, y \in \mathbb{R}\right\}$

3. Solve for  $x$ ,  $\frac{2x+1}{1-x} \geq 1$

A.  $0 \leq x < 1$

B.  $x \leq 0$  or  $x > 1$

C.  $x > 0$  or  $x > 1$


D.  $0 < x \leq 1$

Section I (cont'd)

Marks

4. A particle is oscillating in Simple Harmonic Motion where its position  $x$  metres from a fixed point  $O$  on the same line as its motion after  $t$  seconds is given by  $x = 2 \cos\left(3t + \frac{\pi}{6}\right)$ . What is the maximum speed of the particle?
- A. 2 m/s
  - B. 6 m/s
  - C. 0 m/s
  - D.  $\frac{\pi}{9}$  m/s

5.



$AB$  is a common chord to the circles with centres  $G$  and  $H$ .

$PQ$  is drawn intersecting circle centre  $G$  at  $P$  and  $N$ , intersecting circle centre  $H$  at  $M$  and  $Q$  and intersecting  $AB$  at  $X$  as shown in the diagram.

If  $PM = 18$ ,  $MX = 6$  and  $NQ = 15$  then the length  $NX$  is:

- A. 5
  - B. 4
  - C. 3
  - D. 2
6. The derivative of  $\tan^{-1} \frac{2x}{3}$  is:
- A.  $\frac{1}{3+4x^2}$
  - B.  $\frac{1}{\frac{9}{4}+x^2}$
  - C.  $\frac{6}{9+4x^2}$
  - D.  $\frac{3}{4+x^2}$

Section I (cont'd)

Marks

7. The exact value of  $\sin^{-1}\left(\cos\frac{2\pi}{3}\right)$  is:
- A.  $\frac{\pi}{6}$
  - B.  $-\frac{\pi}{6}$
  - C.  $\frac{\pi}{3}$
  - D.  $-\frac{\pi}{3}$
8. Consider  $(1 + 2x)^n$ . If the ratio of the coefficient of  $x^4$  to the coefficient of  $x^6$  is 5:8 then the value of  $n$  is:
- A. 5
  - B. 6
  - C. 7
  - D. 8
9. The polynomial  $P(x) = x^4 - 2x^3 - 7x^2 + 20x - 12$  has a zero of multiplicity 2 at  $x =$  :
- A. 1
  - B. -3
  - C. 2
  - D. -2
10. A particle moves in a straight line. At time  $t$  seconds, where  $t \geq 0$ , its displacement  $x$  metres from the origin and its velocity  $v$  metres per second are such that  $v = 25 + x^2$ .
- If  $x = 5$  initially, then  $t$  is equal to:
- A.  $25x + \frac{x^3}{3}$
  - B.  $25x + \frac{x^3}{3} + \frac{500}{3}$
  - C.  $\tan^{-1}\left(\frac{x}{5}\right) - \frac{\pi}{4}$
  - D.  $\frac{1}{5}\tan^{-1}\left(\frac{x}{5}\right) - \frac{\pi}{20}$

Section II – Show all working

Question 11 – Start A New Booklet – (15 marks)

Marks

a) (i) Find the derivative of  $\log_e(\cos^2 x)$

1

(ii)  $\int_0^1 \frac{x^2}{x^3 + 1} dx$

1

b) In the expression of  $\left(x^2 + \frac{2}{x}\right)^{10}$  find the coefficient of  $x^2$

1

c) The quadratic polynomial  $ax^2 + bx + 14$  leaves a remainder of  $-12$  when divided by  $(x - 1)$ , and has  $(x + 2)$  as a factor. Find the values of  $a$  and  $b$ .

2

d) Find the acute angle between the lines  $y = 5 - x$  and  $\sqrt{3}y = x + 1$

1

e) (i) Show that the area of an equilateral triangle of side length  $x$  is given by

1

$$A = \frac{\sqrt{3}}{4}x^2$$


(ii) The sides of an equilateral triangle are increasing at the rate of 5 mm/s. At what rate is the area of the triangle increasing at the instant the sides are 10 cm long.

2

Question 11 (cont'd)

Marks

f)



Two circles touch internally at a point  $A$  and the smaller of the two circles passes through  $O$ , the centre of the larger circle.

$AB$  is any chord of the larger circle at  $S$ . The tangents to the larger circle at  $A$  and  $B$  meet at the point  $T$

Prove:

- (i)  $AB$  is bisected at  $S$ . 4
  
- (ii)  $O, S$  and  $T$  are collinear. 2

Question 12 – Start A New Booklet – (15 marks)

Marks

- a) Use the principle of Mathematical Induction to prove that  $7^n + 2$  is divisible by 3 for all positive integers  $n$

2

- b) (i) Show that  $\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$

1

- (ii) Hence or otherwise find the exact value of:

2

$$\int_0^{\frac{\pi}{6}} \sin 4x \cos 2x \, dx$$

- c) (i) Show that  $\frac{d}{dx} (x - \tan^{-1}x) = \frac{x^2}{1+x^2}$

1

- (ii) Hence or otherwise find the exact value of

1

$$\int_0^1 \frac{x^2}{1+x^2} \, dx$$

- d) Given  $A(-2, 3)$  and  $B(4, 7)$  find the coordinates of the point which divides the interval  $AB$  externally in the ratio 3:1

1

- e) If  $\alpha, \beta$  and  $\gamma$  are the roots of  $x^3 - 2x^2 + 4x - 7 = 0$  evaluate  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$

2

Question 12 (cont'd)

Marks

- f) A particle moving in a straight line has an acceleration given by  $\ddot{x} = x^2$  where its displacement is  $x$  metres from the origin. If initially the particle is at rest 2 metres from the origin, find its velocity when it is 4 metres from the origin. 2
- g) The normal at  $P(2ap, ap^2)$  to the parabola  $x^2 = 4ay$  meets the curve again at  $Q(2aq, aq^2)$
- (i) Given that the equation of the normal at  $P$  is  $x + py = ap^3 + 2ap$  1  
show that  $q = -\frac{(2+p^2)}{p}$
- (ii) Find a value for  $p$  so that the lines  $OP$  and  $OQ$  are at right angles, where  $O$  is the origin. 2



Question 13 – Start A New Booklet – (15 marks)

Marks

a) If  $3n^2 - 7n + 5 \equiv An(n - 1) + Bn + C$  find  $A, B$  and  $C$

2

b) Evaluate, leaving your answer in exact form

3

$$\int_{\frac{1}{8}}^{\frac{\sqrt{3}}{8}} \frac{dx}{\sqrt{1 - 16x^2}}$$

c) If  $a$  and  $\beta$  are the roots of  $x^2 + bx + c = 0$ , form the equation, in general form, whose roots are  $\frac{\alpha}{\beta}$  and  $\frac{\beta}{\alpha}$

2

d) A curve is defined by the parametric equations  $x = t - 3$ ,  $y = t^2 - 9$

(i) Find  $\frac{dy}{dx}$  in terms of  $t$

1


(ii) Find the equation of the tangent to the curve at the point where  $t = -3$

2

Question 13 (cont'd)

Marks

e)



$PQ$  is a chord of the parabola  $x^2 = 8y$  passing through the point  $A(0, 1)$  where  $P$  is  $(4p, 2p^2)$  and  $Q$  is  $(4q, 2q^2)$

The tangents to the parabola at  $P$  and  $Q$  meet at the point  $T$ .

$R$  is a point on the chord  $PQ$  with  $RT \perp PQ$

- (i) Write down the equations of the tangents at  $P$  and  $Q$  and hence find the coordinates of  $T$  2

- (ii) Show that the equation of the chord  $PQ$  is given by 1

$$2y = (p + q)x - 4pq$$


- (iii) Show that  $pq = -\frac{1}{2}$  1

- (iv) Find the equation of  $RT$  1

Question 14 – Start A New Booklet – (15 marks)

Marks

a)



A vertical tower  $BT$  of height 18 metres stands with its base  $B$  on horizontal grounds.  $B$  is due North of a fixed point  $P$  and the angle of elevation from  $P$  to the top of the tower  $T$  is  $\frac{\pi}{4}$  radians.  $Q$  is a moving point on the ground due East of  $B$  and the angle of elevation from  $Q$  to  $T$  is  $\theta$  radians where  $0 < \theta < \frac{\pi}{2}$ . The size of the angle  $\theta$  is increasing at a constant rate of 0.02 radians per minute.

(i) Show that  $PQ = 18 \operatorname{cosec} \theta$  2

(ii) Find the rate at which the length  $PQ$  is changing when  $\theta = \frac{\pi}{3}$  2

b) A person hits a ball off the ground with a bat, projecting the ball at a velocity of 50 m/s at an angle of projection  $\theta$  such that  $\tan \theta = \frac{3}{4}$

(i) Taking the origin as the point of projection and  $g = 10 \text{ m/s}^2$  show that  $\dot{x} = 40$  and  $\dot{y} = -10t + 30$  and then find  $x$  and  $y$  in terms of  $t$  3

(ii) A tall building is 100 m from where the ball is hit on horizontal ground. If the ball passes through a small open window in the building find the height of the window above the ground. 2

(iii) Find the velocity and angle that the ball makes with the horizontal as it passes through the window. 2

Question 14 continued on next page

Question 14 (cont'd)

Marks

c) Find the general solution in radians of the equation  $\sin 2x = \cos x$  2

d) By considering the expansion of both sides of the identity

$(1 + x)^{m+n} = (1 + x)^m(1 + x)^n$ , where  $m$  and  $n$  are positive integers, show that

$$\binom{m+n}{3} = \binom{m}{3} + \binom{m}{2}\binom{n}{1} + \binom{m}{1}\binom{n}{2} + \binom{n}{3}$$

Student Number: \_\_\_\_\_ Teacher: \_\_\_\_\_

**Year 12 Mathematics Extension 1 Trial HSC Examination 2012**

**Section I**

**Multiple-choice Answer Sheet – Questions 1 – 10**

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample  $2 + 4 =$  (A) 2      (B) 6      (C) 8      (D) 9  
A       B       C       D

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A       B       C       D

If you change your mind and have crossed out what you consider to be the correct answer, then indicate this by writing the word *correct* and drawing an arrow as follows:

A       B  <sup>correct</sup>      C       D

- 
- |     |   |                       |   |                       |   |                       |   |                       |
|-----|---|-----------------------|---|-----------------------|---|-----------------------|---|-----------------------|
| 1.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 2.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 3.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 4.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 5.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 6.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 7.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 8.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 9.  | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |
| 10. | A | <input type="radio"/> | B | <input type="radio"/> | C | <input type="radio"/> | D | <input type="radio"/> |

Question 11

$$(a) (i) \frac{d \log_e(\cos^2 x)}{dx} = \frac{2 \cos x (-\sin x)}{\cos^2 x} \quad (\cos x \neq 0)$$

$$= -2 \tan x$$

$$(ii) \int_0^1 \frac{x^2}{x^3+1} dx = \frac{1}{3} \int_0^1 \frac{3x^2}{x^3+1} dx$$

$$= \frac{1}{3} [\log_e(x^3+1)]_0^1$$

$$= \frac{1}{3} (\log_e 2 - \log_e 1)$$

$$= \frac{1}{3} \log_e 2$$

$$(b) T_{k+1} = {}^{10}C_k (x^2)^k \left(\frac{2}{x}\right)^{10-k}$$

$$= {}^{10}C_k x^{2k} 2^{10-k} x^{k-10}$$

$$= {}^{10}C_k 2^{10-k} x^{3k-10}$$

If  $3k - 10 = 2$   
 $k = 4$

Coeff of  $x^2 = {}^{10}C_4 \times 2^6$

(c) Let  $P(x) = ax^2 + bx + 14$       ① + ②  $3a = -33$   
 $P(1) = -12$        $a = -11$   
 $a + b + 14 = -12$       Subst in ①  
 $a + b = -26$       ①       $-11 + b = -26$   
 $P(-2) = 0$        $b = -15$   
 $4a - 2b + 14 = 0$   
 $4a - 2b = -14$        $a = -11$        $b = -15$   
 $2a - b = -7$       ②

(d)

$$y = 5 - x$$

$$m_1 = -1$$

$$\sqrt{3}y = x + 1$$

$$m_2 = \frac{1}{\sqrt{3}}$$

Let  $\theta$  be the acute angle between the lines

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$


$$= \left| \frac{-1 - \frac{1}{\sqrt{3}}}{1 + -1 \times \frac{1}{\sqrt{3}}} \right|$$

$$= \left| \frac{-\sqrt{3} - 1}{\sqrt{3} - 1} \right|$$

$$= \frac{\sqrt{3} + 1}{\sqrt{3} - 1}$$

$$\therefore \theta = 75^\circ$$

(e) (i)



$$A = \frac{1}{2} x \times x \sin 60^\circ$$

$$= \frac{1}{2} x^2 \times \frac{\sqrt{3}}{2}$$

$$= \frac{\sqrt{3}}{4} x^2$$

(ii)

$$\frac{dx}{dt} = 0.5 \text{ cm/s}$$

$$\frac{dA}{dt} = \frac{dA}{dx} \frac{dx}{dt}$$

$$= \frac{\sqrt{3}}{2} x \cdot 0.5$$


When  $x = 10$ 

$$\frac{dA}{dt} = \frac{\sqrt{3}}{2} \times 10 \times 0.5$$

$$= \frac{5\sqrt{3}}{2}$$

$\therefore$  Area is increasing at a rate of  $\frac{5\sqrt{3}}{2} \text{ cm}^2/\text{s}$  when

$$x = 10$$



Join OA  
 OA is a radius of larger circle  
 $\angle OAT = 90^\circ$  (angle between radius and tangent at point of contact)

AT is a tangent to smaller circle  
 Since  $\angle TAO = 90^\circ$  AO must pass through centre of smaller circle  
 $\therefore$  AO is a diameter of smaller circle.

Join OS  $\therefore \angle OSA = 90^\circ$  (angle in a semicircle)  
 Join OB

$OB = OA$  (radius of larger circle)  
 $\therefore \triangle OBA$  is isosceles  
 Since  $OS \perp AB$  then OS bisects AB (perp to base of isosceles  $\triangle$  bisects base)  
 ie S is the midpoint of AB

(ii)  $\triangle ABT$  is isosceles  $BT = AT$  (tangents from external point equal in length)  
 Since S is midpt of AB then  $TS \perp AB$

$$\therefore \angle TSO = 90^\circ + 90^\circ = 180^\circ$$

$\therefore$  TSO is a straight angle

ie O, S, T are collinear



## Question 12

(a) Aim: To prove  $7^n + 2$  is divisible by 3  
ie  $7^n + 2 = 3A$  where  $A$  is an integer

$$\text{For } n=1 \quad 7^1 + 2 = 9 = 3 \times 3$$

$\therefore$  Proposition true for  $n=1$

Assume proposition is true for  $n=k$  where  $k$  is a positive integer

$$\text{ie } 7^k + 2 = 3B \text{ where } B \text{ is an integer}$$

Aim to show that proposition is then true for  $n=k+1$

$$\begin{aligned} 7^{k+1} + 2 &= 7 \times 7^k + 2 \\ &= 7 \times (3B - 2) + 2 \quad (\text{by inductive hypothesis}) \\ &= 7 \times 3B - 14 + 2 \\ &= 3 \times 7B - 12 \\ &= 3(7B - 4) \\ &= 3C \quad (C \text{ is an integer since integers} \\ &\quad \text{closed under mult}^n \text{ and subtraction}) \end{aligned}$$

ie proposition is true for  $n=k+1$  if true for  $n=k$ .

Hence by induction proposition is true for all positive integers  $n$ .

$$\begin{aligned} (k) \text{ (i) } \sin(A+B) + \sin(A-B) &= \sin A \cos B + \cos A \sin B \\ &\quad + \sin A \cos B - \cos A \sin B \\ &= 2 \sin A \cos B \end{aligned}$$

$$\begin{aligned} \text{(ii) } \sin 4x \cos 2x &= \frac{1}{2} (\sin(4x+2x) + \sin(4x-2x)) \\ &= \frac{1}{2} (\sin 6x + \sin 2x) \end{aligned}$$

$$\begin{aligned}
\therefore \int_0^{\frac{\pi}{6}} \sin 4x \cos 2x \, dx &= \frac{1}{2} \int_0^{\frac{\pi}{6}} \sin 6x + \sin 2x \, dx \\
&= \frac{1}{2} \left[ -\frac{1}{6} \cos 6x - \frac{1}{2} \cos 2x \right]_0^{\frac{\pi}{6}} \\
&= \left( -\frac{1}{12} \cos \pi - \frac{1}{4} \cos \frac{\pi}{3} \right) - \\
&\quad \left( -\frac{1}{12} \cos 0 - \frac{1}{4} \cos 0 \right) \\
&= \left( -\frac{1}{12} \times -1 - \frac{1}{4} \times \frac{1}{2} \right) - \left( -\frac{1}{12} \times 1 - \frac{1}{4} \times 1 \right) \\
&= \frac{1}{12} - \frac{1}{8} + \frac{1}{12} + \frac{1}{4} \\
&= \frac{7}{24}
\end{aligned}$$

(c) (i)  $\frac{d}{dx} (x - \tan^{-1} x) = 1 - \frac{1}{1+x^2}$

$$\begin{aligned}
&= \frac{1+x^2-1}{1+x^2} \\
&= \frac{x^2}{1+x^2}
\end{aligned}$$

(ii)  $\int_0^1 \frac{x^2}{1+x^2} \, dx = \left[ x - \tan^{-1} x \right]_0^1$

$$\begin{aligned}
&= (1 - \tan^{-1} 1) - (0 - \tan^{-1} 0) \\
&= 1 - \frac{\pi}{4} - 0 \\
&= \frac{4-\pi}{4}
\end{aligned}$$

(d)  $(-2, 3) \quad (4, 7)$   
 $-3:1$

$$\left( \frac{1 \times -2 + -3 \times 4}{-3+1}, \frac{1 \times 3 + -3 \times 7}{-3+1} \right) = \left( \frac{-14}{-2}, \frac{-18}{-2} \right)$$

Point is  $(7, 9)$

(e)

 $x^3 - 2x^2 + 4x - 7 = 0$  has roots  $\alpha, \beta, \gamma$ 

$$\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma} \quad \frac{4}{1} - \frac{-7}{1}$$

$$= \frac{4}{7}$$

(f)

$$\ddot{x} = x^2$$

$$d\left(\frac{1}{2}v^2\right) = x^2 dx$$

$$\frac{1}{2}v^2 = \frac{x^3}{3} + C_1$$

When  $t=0$   $v=0$   $x=2$ 

$$0 = \frac{8}{3} + C_1$$

$$C_1 = -\frac{8}{3}$$

$$\frac{1}{2}v^2 = \frac{x^3}{3} - \frac{8}{3}$$

$$v^2 = \frac{2x^3}{3} - \frac{16}{3}$$

When  $x=4$ 

$$v^2 = \frac{2 \times 64}{3} - \frac{16}{3}$$

$$= \frac{112}{3}$$

$$v = \pm \sqrt{\frac{112}{3}}$$

But  $v \geq 0$ 

$$\therefore v = \sqrt{\frac{112}{3}}$$

OR When  $t=0$   $v=0$   $x=-2$ 

$$0 = -\frac{8}{3} + C_2$$

$$C_2 = \frac{8}{3}$$

$$\frac{1}{2}v^2 = \frac{x^3}{3} + \frac{8}{3}$$

$$v^2 = \frac{2x^3}{3} + \frac{16}{3}$$

Since  $\ddot{x} \geq 0$  ( $\ddot{x} = x^2$ )and  $v=0$  when  $t=0$ 

particle will always move in a positive direction

$$\therefore x \neq -4$$

ie When  $x=4$ 

$$v^2 = \frac{2 \times 64}{3} + \frac{16}{3}$$

$$= \frac{144}{3}$$

$$v = \sqrt{\frac{144}{3}}$$

$$= \sqrt{48} = 4\sqrt{3}$$

(9) (i)  $x + py = ap^3 + 2ap$   
 meets  $x^2 = 4ay$   
 $x + p \cdot \frac{x^2}{4a} = ap^3 + 2ap$

$$\frac{p}{4a} x^2 + x - (ap^3 + 2ap) = 0$$

has roots  $2ap$  and  $2aq$

$$2ap + 2aq = \frac{-1}{p/4a}$$

$$= \frac{-4a}{p}$$

$$2aq = \frac{-4a}{p} - 2ap$$

$$q = \frac{-2}{p} - p$$

$$= \frac{-2 - p^2}{p}$$

$$= \frac{-(2 + p^2)}{p}$$

(ii) Grad OP =  $\frac{ap^2 - 0}{2ap - 0}$

$$= \frac{p}{2}$$

Grad OQ =  $\frac{q}{2}$

if  $OP \perp OQ$  then  $\frac{p}{2} \times \frac{q}{2} = -1$

$$pq = -4$$

$$\therefore q = \frac{-4}{p}$$

$$\therefore \frac{-4}{p} = -\frac{(2+p^2)}{p}$$

$$2+p^2 = 4$$

$$p^2 = 2$$

$$p = \pm\sqrt{2}$$

## Question 13

(a)  $3n^2 - 7n + 5 \equiv An(n-1) + Bn + C$

2 marks

Let  $n=0$   $5 = 0 + 0 + C$   $C=5$

Coeff  $n^2$   $3 = A$   $A=3$

Let  $n=1$   $3 - 7 + 5 = 0 + B + C$

$1 = B + 5$

$B = -4$

(b) 
$$\int_{\frac{1}{8}}^{\frac{\sqrt{3}}{8}} \frac{dx}{\sqrt{1-16x^2}} = \int_{\frac{1}{8}}^{\frac{\sqrt{3}}{8}} \frac{dx}{4\sqrt{\frac{1}{16}-x^2}}$$

$$= \frac{1}{4} \left[ \sin^{-1} \frac{x}{\frac{1}{4}} \right]_{\frac{1}{8}}^{\frac{\sqrt{3}}{8}}$$

$$= \frac{1}{4} \left[ \sin^{-1} 4x \right]_{\frac{1}{8}}^{\frac{\sqrt{3}}{8}}$$

3 marks

$$= \frac{1}{4} \left( \sin^{-1} \frac{\sqrt{3}}{2} - \sin^{-1} \frac{1}{2} \right)$$

$$= \frac{1}{4} \left( \frac{\pi}{3} - \frac{\pi}{6} \right)$$

$$= \frac{\pi}{24}$$

(c)  $x^2 + bx + c = 0$

2 marks

$$2\alpha + \beta = -b$$

equation of quad  
with  $\frac{\alpha}{\beta}$  &  $\frac{\beta}{\alpha}$

$$2\beta = \frac{c}{\alpha}$$

$$\begin{aligned} \frac{\alpha}{\beta} + \frac{\beta}{\alpha} &= \frac{\alpha^2 + \beta^2}{\alpha\beta} \\ &= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} \\ &= \frac{u^2 - 2c}{c} \end{aligned}$$

$$\frac{\alpha}{\beta} - \frac{\beta}{\alpha} = 1$$

$$\text{Eq}^n \text{ is } \left(x - \frac{\alpha}{\beta}\right) \left(x - \frac{\beta}{\alpha}\right) = 0$$

$$x^2 - \left(\frac{\alpha}{\beta} + \frac{\beta}{\alpha}\right)x + \frac{\alpha}{\beta} \cdot \frac{\beta}{\alpha} = 0$$

$$x^2 - \frac{(u^2 - 2c)}{c}x + 1 = 0$$

$$cx^2 - (u^2 - 2c)x + c = 0$$

(a) Eq<sup>n</sup> of tangents at P and Q

$$y = px - 2p^2 \quad (1)$$

$$y = qx - 2q^2 \quad (2)$$

Subst (1) in (2)

$$px - 2p^2 = qx - 2q^2$$

$$(p - q)x = 2(p^2 - q^2)$$

$$x = \frac{2(p - q)(p + q)}{p - q} \quad (p \neq q)$$

$$= 2(p + q)$$

$$y = p \cdot 2(p + q) - 2p^2$$

$$= 2pq$$

$\therefore T$  is point  $(2(p + q), 2pq)$

2 marks.

Write equations of  
tangent  
with  $p, q$

$$\begin{aligned}
 \text{(ii) Grad PQ} &= \frac{2p^2 - 2q^2}{4p - 4q} \\
 &= \frac{2(p-q)(p+q)}{4(p-q)} \\
 &= \frac{p+q}{2}
 \end{aligned}$$

Eq<sup>n</sup> of chord PQ is

$$y - 2p^2 = \frac{p+q}{2} (x - 4p)$$

$$= \frac{p+q}{2} \cdot x - 2p(p+q) \quad \text{1 mark}$$

$$= \frac{p+q}{2} x - 2p^2 - 2pq$$

$$y = \frac{p+q}{2} x - 2pq$$

$$2y = (p+q)x - 4pq$$

(iii) Since PQ passes through A(0,1)

$$2 = 0 - 4pq$$

$$pq = -\frac{1}{2}$$

1 mark

(iv) RT  $\perp$  PQ

$$\therefore \text{Grad RT} = -\frac{2}{p+q}$$

1 mark  
equation RT

$\therefore$  Eq<sup>n</sup> of RT is

$$y - 2pq = -\frac{2}{p+q} (x - 2(p+q))$$

$$y - 2 \times -\frac{1}{2} = -\frac{2}{p+q} x + 4$$

$$y = -\frac{2}{p+q} x + 3$$



$$(a) (i) \quad x = t - 3 \quad y = t^2 - 9$$

1 mark

$$\frac{dx}{dt} = 1$$

$$\frac{dy}{dt} = 2t$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$= \frac{2t}{1}$$

$$(ii) \text{ When } t = -3$$

2 marks

$$x = -6$$

$$y = 0$$


$$\frac{dy}{dx} = -6$$

Eq<sup>n</sup> of tangent is

$$y - 0 = -6(x - (-6))$$

$$y = -6x - 36$$

# Question 14



$$(i) \quad \frac{BQ}{18} = \cot \theta$$

$$BQ = 18 \cot \theta$$

$$\frac{BP}{18} = \tan \frac{\pi}{4}$$

$$= 1$$

$$BP = 18$$

$$\begin{aligned} \text{In } \triangle PBQ \quad PQ^2 &= BP^2 + BQ^2 \\ &= 18^2 + 18^2 \cot^2 \theta \\ &= 18^2 (1 + \cot^2 \theta) \\ &= 18^2 \operatorname{cosec}^2 \theta \end{aligned}$$

$$PQ = 18 \operatorname{cosec} \theta$$

$$(ii) \quad \frac{dPQ}{dt} = \frac{dPQ}{d\theta} \cdot \frac{d\theta}{dt}$$

$$= -18 \operatorname{cosec} \theta \cot \theta \cdot \frac{d\theta}{dt}$$



$$= -18 \operatorname{cosec} \theta \cot \theta \cdot 0.02$$

$$= -0.36 \times \operatorname{cosec} \frac{\pi}{3} \cdot \cot \frac{\pi}{3}$$

$$= -0.36 \times \frac{2}{\sqrt{3}} \times \frac{1}{\sqrt{3}}$$

$$= -0.24$$

(b) (i)



$$\ddot{x} = 0$$

$$\dot{x} = c_1$$

$$\text{When } t=0 \quad \dot{x} = 50 \cos \theta \\ = 40$$

$$\therefore c_1 = 40$$

$$\dot{x} = 40$$

$$x = 40t + c_3$$

$$\text{When } t=0 \quad x=0 \\ \therefore c_3 = 0$$

$$x = 40t$$

$$\ddot{y} = -10$$

$$\dot{y} = -10t + c_2$$

$$\text{When } t=0 \quad \dot{y} = 50 \sin \theta \\ = 30$$

$$\therefore c_2 = 30$$

$$\dot{y} = -10t + 30$$

$$y = -5t^2 + 30t + c_4$$

$$\text{When } t=0 \quad y=0 \\ \therefore c_4 = 0$$

$$y = -5t^2 + 30t$$

(ii) If  $x = 100$  then

$$100 = 40t$$

$$t = \frac{5}{2}$$


$$y = -5 \times \left(\frac{5}{2}\right)^2 + 30 \times \frac{5}{2}$$

$$= \frac{175}{4}$$

$\therefore$  Height of window is  $43.75 \text{ m}$

(iii) When  $t = \frac{5}{2}$   $\dot{x} = 40$

$$\dot{y} = -10 \times \frac{5}{2} + 30 = 5$$



$$v^2 = 40^2 + 5^2$$

$$= 1600 + 25$$

$$= 1625$$

$$v = \sqrt{1625}$$

$$= 5\sqrt{65}$$

$$\tan \theta = \frac{5}{40}$$

$$= \frac{1}{8}$$

$$\theta = \tan^{-1}\left(\frac{1}{8}\right)$$

$$= 7^\circ 8'$$

Velocity is  $5\sqrt{65}$  m/s and angle ball's path makes with the horizontal is  $7^\circ 8'$

(c)  $\sin 2x = \cos x$

$$2\sin x \cos x - \cos x = 0$$

$$\cos x (2\sin x - 1) = 0$$

$$\cos x = 0 \quad \text{or} \quad \sin x = \frac{1}{2}$$

$$x = \frac{\pm \pi}{2}, \frac{\pm 3\pi}{2}, \frac{\pm 5\pi}{2}, \dots \quad x = \frac{\pi}{6} + 2k\pi, \pi - \frac{\pi}{6} + 2k\pi$$

$$x = \frac{(2k+1)\pi}{2}, \frac{\pi}{6} + 2k\pi, \frac{5\pi}{6} + 2k\pi \quad (k \in \mathbb{Z})$$

(d)  $(1+x)^{m+n} = (1+x)^m (1+x)^n$

On LHS coeff  $x^3 = \binom{m+n}{3}$

$$\text{RHS} = \left( 1 + \binom{m}{1}x + \binom{m}{2}x^2 + \binom{m}{3}x^3 + \dots \right)$$

$$\times \left( 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots \right)$$

$$\text{Term in } x^3 = 1 \times \binom{n}{3}x^3 + \binom{m}{1}x \times \binom{n}{2}x^2 + \binom{m}{2}x \times \binom{n}{1}x$$

$$+ 1 \times \binom{m}{3}x^3$$

$$\therefore \text{Coeff } x^3 = \binom{n}{3} + \binom{m}{1} \times \binom{n}{2} + \binom{m}{2} \times \binom{n}{1} + 1 \times \binom{m}{3}$$

$$= \binom{m+n}{3}$$

