



2009
TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics

Extension 2

General Instructions

- Reading Time- 5 minutes
- Working Time – 3 hours
- Write using a blue or black pen
- Approved calculators may be used
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown for every question.
- Use a separate answer booklet for each question

Total marks - 120


- Attempt Questions 1-8
- All questions are of equal value

Question 1 – 15 marks – Use a separate writing booklet

- | | Marks |
|--|--------------|
| (a) Find $\int \frac{\cos \theta}{\sin^3 \theta} d\theta$ | 2 |
| (b) Use completing the square to find $\int \frac{1}{\sqrt{x^2 + 4x + 13}} dx$ | 2 |
| (c) Use integration by parts to find $\int_0^1 \tan^{-1} x dx$ | 3 |
| (d) Find real numbers A and B so that | 2 |
| $\frac{4x^2 - 2x + 9}{(x+1)(x^2 + 4)} \equiv \frac{A}{x+1} + \frac{Bx - 3}{x^2 + 4}$ | |
| Hence find $\int \frac{4x^2 - 2x + 9}{(x+1)(x^2 + 4)} dx$ | 2 |
| (e) Use the substitution $x = 2 \sin \theta$ to find $\int_0^1 \frac{1}{(4 - x^2)^{\frac{3}{2}}} dx$ | 4 |

Question 2 – 15 marks – Use a separate writing booklet

- | | Marks |
|---|--------------|
| (a) Let $z = 2 + i$ | |
| (i) Find z^3 in the form $a + ib$ | 1 |
| (ii) Find $\frac{5}{z}$ in the form $a + ib$ | 1 |
| (b) Find all pairs of real numbers x and y such that $(x + iy)^2 = 3 + 4i$. | 2 |
| (c) Let $\alpha = 1 - i$ | |
| (i) Express α in modulus-argument form. | 2 |
| (ii) Hence evaluate α^{10} in the form $a + ib$ | 2 |
| (d) On separate Argand diagrams, sketch the region satisfying the following inequalities: | |
| (i) $1 \leq z \leq 2$ | 1 |
| (ii) $\frac{\pi}{4} \leq \arg(z) \leq \frac{3\pi}{4}$ | 1 |
| (iii) $0 \leq \operatorname{Re}(z) \leq 3$ and $1 \leq \operatorname{Im}(z) \leq 2$ | 2 |



On the Argand diagram shown, $ABCD$ is a square.

The points A and B represent complex numbers α and β respectively.

Hence AB represents the complex number $\beta - \alpha$.

Find, in terms of α and β ,

- | | |
|---|----------|
| (i) the complex number represented by AD | 1 |
| (ii) the complex number represented by BD | 1 |
| (iii) the complex number that is represented by the point C | 1 |


Question 3 – 15 marks . Use a separate writing booklet

Marks

- (a) Find the gradient of the tangent to the curve with equation $x^2 - 3xy + y^2 = 5$ at the point (4,1) on the curve.

2

(b)



The diagram shows the graph of $y = f(x)$


Make separate, one-third page sketches of the following:

- (i) $y = f(|x|)$ 1
- (ii) $y = \frac{1}{f(x)}$ 2
- (iii) $y = (f(x))^2$ 2
- (iv) $y = \sqrt{f(x)}$ 2
- (v) $y = 2^{f(x)}$ 2
- (c) Consider the graph of the function $y = \frac{x^2 + 1}{x^2 - 9}$.
- (i) Write down the equations of the vertical and horizontal asymptotes. 2
- (ii) Make a neat sketch of the graph of $y = \frac{x^2 + 1}{x^2 - 9}$ clearly showing the asymptotes and any intercepts with the coordinate axes. 2

Question 4 – 15 marks . Use a separate writing booklet

Marks

(a)



The region bound by $y = (x - 4)^2$ and $y = 4$ is rotated about the y -axis to form a solid.


(i) Use the method of cylindrical shells to form an integral whose value will give the volume of the solid

3

(ii) Evaluate this integral to find the volume of the solid.

2

(b)



The base of a solid is the region bound by $y = 4 - x^2$ and the x -axis. Vertical cross-sections parallel to the x -axis are equilateral triangles. A typical cross-section ABC is shown.

(i) Show that the area of an equilateral triangle with side s units is

1

given by $A = \frac{\sqrt{3}}{4} s^2$


(ii) Form an integral whose value will give the volume of the solid and evaluate this integral to find the volume.

4

Question 4 (Continued)

Marks

(c)



A solid is formed by rotating the circle $x^2 + y^2 = 1$ about the line $x = 4$.
Any cross-section perpendicular to the line $x = 4$ will be an annulus.

- (i) Consider the annulus formed when the chord AB, at $y = h$, is rotated about the line $x = 4$.

2

Show that the area of this annulus is given by $A = 16\pi\sqrt{1-h^2}$

- (ii) Form an integral whose value will give the volume of the solid and evaluate it to find the volume of the solid.

3

Question 5 – 15 marks . Use a separate writing booklet


- | | Marks |
|---|--------------|
| (a) The polynomial $P(x) = 2x^4 + 9x^3 + 6x^2 - 20x - 24$ has a triple zero.
Find all the zeros of $P(x)$. | 3 |
| (b) The cubic equation $x^3 - x^2 + 4x - 2 = 0$ has roots α , β and γ . | |
| (i) Find the cubic polynomial with integer coefficients whose roots are α^2 , β^2 and γ^2 . | 3 |
| (ii) Find the value $\alpha^2 + \beta^2 + \gamma^2$ | 1 |
| (iii) Find the value of $\alpha^3 + \beta^3 + \gamma^3$ | 1 |
| (c) It is given that $1 + 2i$ is a root of $x^3 + ax^2 + bx + 10 = 0$ where a and b are real. | |
| (i) Find all the roots of $x^3 + ax^2 + bx + 10 = 0$. | 2 |
| (ii) Hence express $x^3 + ax^2 + bx + 10$ as a product of linear and quadratic factors with real coefficients. | 1 |
| (d) Let $z = \cos \theta + i \sin \theta$ | |
| (i) Find the roots of $z^5 = -1$, expressing the complex roots in modulus-argument form. | 2 |
| (ii) Hence show that $\cos \frac{\pi}{5} + \cos \frac{3\pi}{5} = \frac{1}{2}$ | 2 |

Question 6 – 15 marks . Use a separate writing booklet

Marks

- (a) Consider the ellipse whose equation is $\frac{x^2}{9} + \frac{y^2}{4} = 1$.
- (i) Find the eccentricity of the ellipse. 1
 - (ii) Write down the coordinates of the foci and the equations of the directrices. 2
 - (iii) Make a neat sketch of the ellipse, clearly showing the foci and the directrices. 2
 - (iv) Let P be any point on the ellipse. Show that $PS + PS' = 6$, where S and S' are the foci of the ellipse. 2

(b)




$P(5p, \frac{5}{p})$, $p > 0$ and $Q(5q, \frac{5}{q})$, $q > 0$ are two points on the hyperbola $xy = 25$.

- (i) Show that the equation of the chord PQ is $x + pqy = 5(p + q)$ 2
- (ii) Show that the equation of the tangent at P is $x + p^2y = 10p$ 2
- (iii) The tangents at P and Q intersect at R. Show that the coordinates of R are $(\frac{10pq}{p+q}, \frac{10}{p+q})$ 2
- (iv) PQ passes through the point T(0,15). Find the equation of the locus of R. 2

Question 7 – 15 marks . Use a separate writing booklet

Marks

(a)



A body of mass m kg is attached by a light, inextensible string of length ℓ metres to the vertex of a smooth, inverted cone whose semi-vertical angle is θ° . The body remains in contact with the surface of the cone and rotates as a conical pendulum with angular velocity ω radians per second and radius r metres.. The forces acting on the body are tension in the string (T), the normal reaction of the cone (N) and the gravitational force mg .

(i) Copy the diagram and show the forces that are acting on the body. 1

(ii) By resolving forces vertically and horizontally, show that 2

$$T \cos \theta + N \sin \theta = mg$$

$$\text{and } T \sin \theta - N \cos \theta = mr\omega^2$$

(iii) Show that $T = mg \cos \theta + mr\omega^2 \sin \theta$ and find a similar expression for N . 2

(iv) The angular velocity is increased until the body is about to lose contact with the cone. Find an expression for this value of ω in terms of g , r and θ . 2

Question 7 (continued)

Marks

(b) A body of mass m kg is projected vertically under the influence of gravity in a medium whose resistance is mkv^2 where v is its velocity and k is a constant. Its initial velocity is U metres per second.

- (i) Make a neat sketch showing the forces acting on the body as it is moving up. 1
- (ii) Show that its acceleration \ddot{x} is given by $\ddot{x} = -(g + kv^2)$ 1
- (iii) Show that the displacement of the body is given by $x = \frac{1}{2k} \log_e \left(\frac{g + kU^2}{g + kv^2} \right)$ 3
- (iv) Find the maximum height reached by the body. 1
- (v) The body now starts falling back towards the ground. 1
Make a neat sketch showing the forces acting on the body and hence write an equation for the acceleration of the body.
- (vi) Determine the terminal velocity of the body as it falls. 1


Question 8 – 15 marks . Use a separate writing booklet

Marks

- (a) Use mathematical induction to prove that $\sin(x + n\pi) = (-1)^n \sin x$
where n an integer ≥ 1

3

- (b) Two circles intersect at A and B and a common tangent touches them at P and Q as shown.



A chord PR is drawn parallel to QA.

RA is produced to meet the other circle at S.

Copy the diagram onto your own paper.

- (i) Explain why $\angle PQA = \angle QSA$.
- (ii) Prove that PQSR is a cyclic quadrilateral
- (iii) Hence prove that PA is parallel to QS

1

3

2

Question 8 (continued)

Marks

(c) Let $I_n = \int_0^{\frac{\pi}{2}} \cos^{2n} x \, dx$ for $n = 0, 1, 2, 3, \dots$

(i) By writing $\int_0^{\frac{\pi}{2}} \cos^{2n} x \, dx$ as $\int_0^{\frac{\pi}{2}} \cos^{2n-1} x \cos x \, dx$ and using integration **4**

by parts, show that $I_n = \frac{2n-1}{2n} I_{n-1}$ for $n = 1, 2, 3, \dots$

(ii) Hence evaluate $\int_0^{\frac{\pi}{2}} \cos^6 x \, dx$ **2**

END OF EXAMINATION